

O. V. Bagro, S. A. Kruglyak

**A MAJORIZATION RELATION FOR A CERTAIN CLASS
OF *-QUIVERS WITH AN ORTHOGONALITY
CONDITION**

Abstract. In [1, 2, 3], *-algebras and *-categories over the field \mathbb{C} of complex numbers were quasi-ordered with respect to the complexity of the structure of their *-representations with a majorization relation \succ . A notion of *-wildness was also introduced there for an algebra (a category) if the algebra majorizes the *-algebra $C^*(\mathcal{F}_2)$. In this paper, we discuss some methods for proving that an algebra is *-wild and obtain criteria for certain “standard” *-categories (ensembles with an orthogonality condition) to be *-wild.

Keywords: algebras, categories and functors, representations.

Mathematics Subject Classification: 18B30.

1. INTRODUCTION

A majorization relation \succ for *-algebras and *-categories was introduced in [1, 2, 3] as follows: for categories \mathcal{K}_1 and \mathcal{K}_2 , the majorization $\mathcal{K}_1 \succ \mathcal{K}_2$ means that the problem of classifying *-representations of the category \mathcal{K}_2 over the category of Hilbert spaces is contained in that for the category \mathcal{K}_1 .

Let Q be a *-quiver with relations. Let it generate a *-category $\mathcal{K} = \mathcal{K}(Q)$ over the field \mathbb{C} of complex numbers, see [4, 5]. Let $\text{Rep } \mathcal{K}$ ($\text{Rep } Q$) be the category of *-representations of the *-category \mathcal{K} (*-quiver Q), see Section 2 for the definitions.

Let $\mathcal{F}_2 = \mathbb{C} \langle u_1, u_2 \mid u_i u_i^* = u_i^* u_i = e, i = 1, 2 \rangle$. A *-category \mathcal{K} is said to be *-wild [1, 2, 3] if $\mathcal{K} \succ \mathcal{F}_2$. The wildness indicates that the classification problem is extremely complex; in a certain sense, \mathcal{F}_2 majorizes “all other” *-algebras and *-categories, see [6].

An involution quiver Q will be called an *ensemble* if the sets of its vertices and edges can be written as $Q_v = Q_v^{(1)} \cup Q_v^{(2)}$ and $Q_a = Q_a^{(1)} \cup Q_a^{(2)}$, respectively. Here

the subsets that enter in the unions can have a nonempty intersection and even coincide. They are defined as follows. For every pair of vertices in the sets $Q_v^{(1)}$ and $Q_v^{(2)}$, there is precisely one arrow in $Q_a^{(1)}$ going from the vertex in the set $Q_v^{(1)}$ to the vertex in the set $Q_v^{(2)}$. The same must also be true for each pair of vertices in $Q_v^{(2)}$, $Q_v^{(1)}$, and the arrows in $Q_a^{(2)}$. Hence, if $\alpha \in Q_a^{(1)}$, then $\alpha^* \in Q_a^{(2)}$, and vice versa. If $Q_v^{(1)} = \{a_1, a_2, \dots, a_n\}$, $Q_v^{(2)} = \{b_1, b_2, \dots, b_m\}$ and $\alpha_{b_i a_j} : a_j \rightarrow b_i$, $\alpha_{b_i a_j} \in Q_a^{(1)}$, the ensemble can be given by the two matrices

$$\mathcal{A} = \begin{array}{c} \\ b_1 \\ b_2 \\ \vdots \\ b_m \end{array} \begin{array}{cccc} a_1 & a_2 & \cdots & a_n \\ \alpha_{b_1 a_1} & \alpha_{b_1 a_2} & \cdots & \alpha_{b_1 a_n} \\ \alpha_{b_2 a_1} & \alpha_{b_2 a_2} & \cdots & \alpha_{b_2 a_n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{b_m a_1} & \alpha_{b_m a_2} & \cdots & \alpha_{b_m a_n} \end{array} \quad \text{and} \quad \mathcal{A}^* = \begin{array}{c} \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{array} \begin{array}{cccc} b_1 & b_2 & \cdots & b_m \\ \alpha_{b_1 a_1}^* & \alpha_{b_2 a_1}^* & \cdots & \alpha_{b_m a_1}^* \\ \alpha_{b_1 a_2}^* & \alpha_{b_2 a_2}^* & \cdots & \alpha_{b_m a_2}^* \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{b_1 a_n}^* & \alpha_{b_2 a_n}^* & \cdots & \alpha_{b_m a_n}^* \end{array}$$

An ensemble Q given by the matrices \mathcal{A} and \mathcal{A}^* and the relation

$$\mathcal{A}^* \mathcal{A} = \begin{array}{c} \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{array} \begin{array}{cccc} a_1 & a_2 & \cdots & a_n \\ \varepsilon_{a_1} & & & \\ & \varepsilon_{a_2} & & 0 \\ & 0 & \ddots & \\ & & & \varepsilon_{a_n} \end{array}$$

will be called an ensemble of dimension $m \times n$ with the orthogonality condition

$$\alpha_{1i}^* \alpha_{1j} + \alpha_{2i}^* \alpha_{2j} + \cdots + \alpha_{mi}^* \alpha_{mj} = \begin{cases} 0, & \text{if } i \neq j, \\ \varepsilon_{a_i}, & \text{if } i = j \end{cases}$$

defined in the vertices a_1, a_2, \dots, a_n . This ensemble will be denoted by $Q_{m \times n^\perp}$.

In this paper, we find conditions for majorization of the $*$ -categories, $\mathcal{K}(Q_{k \times l^\perp}) \succ \mathcal{K}(Q_{m \times n^\perp})$ (Theorem 2) and prove a criterion for the $*$ -category $\mathcal{K}(Q_{m \times n^\perp})$ to be $*$ -wild (Theorem 3). As a tool for proving statements on the majorization, we construct a quiver $Q'_{\alpha, x}$ that is derived with respect to the arrow α at the point $x = (x_1, x_2, \dots, x_n)$, $x_i \in \mathbb{R}$, and prove Theorem 1 related to this construction.

2. CATEGORIES WITH AN INVOLUTION AND A MAJORIZATION RELATION

Let \mathcal{K} be a category with an involution $*$ over the field \mathbb{C} of complex numbers [4, 5] such that $a^* = a$ for every $a \in \text{Ob} \mathcal{K}$. Hence, for every morphism $\alpha : a \rightarrow b$ there is a morphism $\alpha^* : b \rightarrow a$ such that 1) $\alpha^{**} = \alpha$, 2) $(\alpha\beta)^* = \beta^* \alpha^*$, 3) $(z_1 \alpha_1 + z_2 \alpha_2)^* = \bar{z}_1 \alpha_1^* + \bar{z}_2 \alpha_2^*$, $z_1, z_2 \in \mathbb{C}$. We assume that there is a zero object in \mathcal{K} . In the sequel, we call such categories $*$ -categories.

Together with the category \mathcal{K} , we will also consider, as a system of generators, an involution quiver (*-quiver) Q [4, 5] such that for each vertex a , $a^* = a$, and for each arrow $\alpha: a \rightarrow b$ there corresponds an arrow $\alpha^*: b \rightarrow a$ such that $\alpha^{**} = \alpha$. The category \mathcal{K} is obtained from the category of paths of Q by factorization. The category \mathcal{K} will be called finitely generated if it can be defined by a finite *-quiver and a finite set of relations (linear combinations of certain paths are set to zero). A morphism $\varphi: a \rightarrow b$ in the category \mathcal{K} is called an isomorphism if there exists a morphism $\varphi^{-1}: b \rightarrow a$ such that $\varphi^{-1}\varphi = \varepsilon_a$ and $\varphi\varphi^{-1} = \varepsilon_b$. An isomorphism φ is called a congruence if $\varphi^{-1} = \varphi^*$. A representation of the category \mathcal{K} is an involutive functor π over the field \mathbb{C} , compatible with the involution, from the category \mathcal{K} to the category \mathcal{H} of Hilbert spaces, where the objects are separable Hilbert spaces and morphisms are bounded linear operators from one space to another. The involution on the objects of \mathcal{H} is an identity, whereas, it is the adjoint operator on a morphism.

Representations of the category \mathcal{K} themselves form a *-category $\text{Rep } \mathcal{K}$ whose objects are involutive functors π (*-representations) and morphisms are families of morphisms of the category \mathcal{H} which intertwine these functors (natural transformations of the functors). Two representations are called equivalent, if they are isomorphic in the category of representations, and are unitary isomorphic, if they are congruent in the category of representations. As is known, see for example [4], that if two *-representations are equivalent, then they are unitary equivalent.

Let us define a *-category of matrices $\mathcal{M}(\mathcal{K})$ over a category \mathcal{K} . Its objects are ordered collections (a_1, a_2, \dots, a_n) of objects of the category \mathcal{K} (we do not exclude the possibility of repetitions of a_i). A morphism from (a_1, a_2, \dots, a_n) into (b_1, b_2, \dots, b_m) is a matrix of dimension $m \times n$ that will be written as

$$\mathcal{A} = \begin{matrix} & a_1 & a_2 & \cdots & a_n \\ \begin{matrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{matrix} & \begin{matrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{matrix} \end{matrix}, \quad \mathcal{A}^* = \begin{matrix} & b_1 & b_2 & \cdots & b_m \\ \begin{matrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{matrix} & \begin{matrix} \alpha_{11}^* & \alpha_{21}^* & \cdots & \alpha_{m1}^* \\ \alpha_{12}^* & \alpha_{22}^* & \cdots & \alpha_{m2}^* \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{1n}^* & \alpha_{2n}^* & \cdots & \alpha_{mn}^* \end{matrix} \end{matrix},$$

$\alpha_{ij} \in \text{Hom}_{\mathcal{K}}(a_j, a_i)$. The composition and addition of morphisms in the category $\mathcal{M}(\mathcal{K})$ is performed by using the usual matrix multiplication and addition.

Let us give a definition of an enveloping *-category, see [1, 2, 3].

Definition 1. Let \mathcal{K} and $\tilde{\mathcal{K}}$ be *-categories. A pair $(\tilde{\mathcal{K}}, \Phi: \mathcal{K} \rightarrow \tilde{\mathcal{K}})$, where Φ is an involutive functor from the *-category \mathcal{K} into the *-category $\tilde{\mathcal{K}}$, is called an enveloping category of the category \mathcal{K} if for any *-representation $\pi: \mathcal{K} \rightarrow \mathcal{H}$ there exists a unique representation $\tilde{\pi}: \tilde{\mathcal{K}} \rightarrow \mathcal{H}$ such that the diagram

$$\begin{array}{ccc}
\tilde{\mathcal{K}} & & \\
\uparrow \Phi & \searrow \tilde{\pi} & \\
\mathcal{K} & \xrightarrow{\pi} & \mathcal{H}
\end{array}$$

is commutative and any morphism in the category $\text{Rep } \mathcal{K}$ that intertwines two representations π_1 and π_2 in the category \mathcal{K} intertwines the representations $\tilde{\pi}_1$ and $\tilde{\pi}_2$ in the category $\tilde{\mathcal{K}}$.

Let $\mathcal{M}(\mathcal{K})$ be the category of matrices over a $*$ -category \mathcal{K} , $(\widetilde{\mathcal{M}(\mathcal{K})}, \Phi: \mathcal{M}(\mathcal{K}) \rightarrow \widetilde{\mathcal{M}(\mathcal{K})})$ its enveloping category. Any $*$ -representation $\pi: \mathcal{K} \rightarrow \mathcal{H}$ induces a representation $\Pi: \mathcal{M}(\mathcal{K}) \rightarrow \mathcal{H}$, hence, a representation $\tilde{\Pi}: \widetilde{\mathcal{M}(\mathcal{K})} \rightarrow \mathcal{H}$. If $\Psi: \mathcal{L} \rightarrow \widetilde{\mathcal{M}(\mathcal{K})}$ is an involutive functor. Then one can naturally construct the functor

$$\mathbf{F}_\Psi: \text{Rep } \mathcal{K} \rightarrow \text{Rep } \mathcal{L}.$$

By the definition, $\mathbf{F}_\Psi(\pi) = \tilde{\Pi} \circ \Psi$ and, if $\mathcal{C} = (\mathcal{C}_a)_{a \in \text{Ob } \mathcal{K}}$ is a morphism of representations in $\text{Rep } \mathcal{K}$, $\mathcal{C}_a: \pi(a) \rightarrow \pi_1(a)$, $(a_1, a_2, \dots, a_k) \in \text{Ob } \mathcal{M}(\mathcal{K})$, then [3]

$$\mathbf{F}_\Psi(\mathcal{C}) = \text{diag}(\mathcal{C}_{a_1}, \mathcal{C}_{a_2}, \dots, \mathcal{C}_{a_k}) \in \text{Hom}_{\mathcal{H}}(\pi(a_1) \oplus \dots \oplus \pi(a_k), \pi_1(a_1) \oplus \dots \oplus \pi_1(a_k)).$$

Definition 2. We will say that a $*$ -category \mathcal{L} directly majorizes a $*$ -category \mathcal{K} and denote this by $\mathcal{L} \triangleright \mathcal{K}$ if the category of matrices $\mathcal{M}(\mathcal{K})$ contains the enveloping category $(\widetilde{\mathcal{M}(\mathcal{K})}, \mathcal{F})$ and a $*$ -functor $\Psi: \mathcal{L} \rightarrow \widetilde{\mathcal{M}(\mathcal{K})}$ such that the functor $\mathbf{F}_\Psi: \text{Rep } \mathcal{K} \rightarrow \text{Rep } \mathcal{L}$, which is univalent by the definition, is full. A $*$ -category \mathcal{L} majorizes a $*$ -category \mathcal{K} , denoted by $\mathcal{L} \succ \mathcal{K}$, if there exist $*$ -categories $\mathcal{K}_1, \mathcal{K}_2, \dots, \mathcal{K}_n$ such that $\mathcal{L} \equiv \mathcal{K}_0 \triangleright \mathcal{K}_1 \triangleright \dots \triangleright \mathcal{K}_n \triangleright \mathcal{K}_{n+1} \equiv \mathcal{K}$. We will say that a $*$ -quiver Q_1 majorizes a $*$ -quiver Q_2 and denote this by $Q_1 \succ Q_2$, if $\mathcal{K}(Q_1) \succ \mathcal{K}(Q_2)$ for the corresponding categories.

The majorization relation is a quasi-order on the class of $*$ -categories and $*$ -quivers.

As a complexity reference for the problem of classifying $*$ -representations of $*$ -algebras and $*$ -categories, we take the classification problem for pairs of self-adjoint operators up to the unitary equivalence [6] or, which is actually the same thing, that for pairs of unitary operators [1, 2, 3].

Definition 3 ([3]). A $*$ -category $\mathcal{K}(Q)$ (a $*$ -quiver Q) is called $*$ -wild, if $\mathcal{K}(Q) \succ \mathcal{F}_2$, where $\mathcal{F}_2 = \mathbb{C} \langle u_1, u_2 \mid u_i u_i^* = u_i^* u_i = e, i = 1, 2 \rangle$.

It is clear that if \mathcal{K} is a $*$ -wild category and $\mathcal{K}_1 \succ \mathcal{K}$, then \mathcal{K}_1 is also $*$ -wild. So, it is useful to have a sufficiently large class of “standard” $*$ -wild algebras (categories). They include, for example, the $*$ -algebra generated by 3 orthogonal projections two of which are mutually orthogonal [1, 2, 6]. In this paper, we construct other “standard” $*$ -wild categories.

3. DERIVED *-QUIVERS AND MAJORIZATION

Let $\alpha \in Q_a$, $\alpha: a \rightarrow b$ and, if $a = b$, then $\alpha = \alpha^*$. Let $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $x_i \neq x_j$ for $i \neq j$ and, in addition, $x_i \geq 0$ if $a \neq b$. If $0 \in \{x_1, x_2, \dots, x_n\}$, we assume that $x_n = 0$.

For a quiver Q , let us define a quiver $Q'_{\alpha, x}$, *derived with respect to the arrow α at the point x* . To construct $Q'_{\alpha, x}$, we do the following.

1. Delete from the quiver Q the arrows α and α^* and replace the vertices a and b with the collections of vertices a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n , gluing the vertices a_i and b_i if $x_i \neq 0$.
2. Each self-adjoint arrow $\beta: a \rightarrow a$ that does not coincide with α is replaced with the ensemble

$$\mathcal{B} = \begin{array}{c} a_1 \\ a_2 \\ \vdots \\ a_n \end{array} \begin{array}{c} a_1 \quad a_2 \quad \cdots \quad a_n \\ \hline \beta_{a_1 a_1} \quad \beta_{a_1 a_2} \quad \cdots \quad \beta_{a_1 a_n} \\ \beta_{a_2 a_1} \quad \beta_{a_2 a_2} \quad \cdots \quad \beta_{a_2 a_n} \\ \vdots \quad \vdots \quad \ddots \quad \vdots \\ \beta_{a_n a_1} \quad \beta_{a_n a_2} \quad \cdots \quad \beta_{a_n a_n} \end{array} = \mathcal{B}^*.$$

The same is done for each self-adjoint arrow $\beta: b \rightarrow b$.

3. Each pair of not self-adjoint arrows $\gamma: a \rightarrow b$ and $\gamma^*: b \rightarrow a$ is replaced with the ensemble

$$\mathcal{T} = \begin{array}{c} a_1 \\ \vdots \\ a_{n-1} \\ b_n \end{array} \begin{array}{c} a_1 \quad \cdots \quad a_{n-1} \quad a_n \\ \hline \gamma_{a_1 a_1} \quad \cdots \quad \gamma_{a_1 a_{n-1}} \quad \gamma_{a_1 a_n} \\ \vdots \quad \ddots \quad \vdots \quad \vdots \\ \gamma_{a_{n-1} a_1} \quad \cdots \quad \gamma_{a_{n-1} a_{n-1}} \quad \gamma_{a_{n-1} a_n} \\ \gamma_{b_n a_1} \quad \cdots \quad \gamma_{b_n a_{n-1}} \quad \gamma_{b_n a_n} \end{array},$$

$$\mathcal{T}^* = \begin{array}{c} a_1 \\ \vdots \\ a_{n-1} \\ a_n \end{array} \begin{array}{c} a_1 \quad \cdots \quad a_{n-1} \quad b_n \\ \hline \gamma_{a_1 a_1}^* \quad \cdots \quad \gamma_{a_{n-1} a_1}^* \quad \gamma_{b_n a_1}^* \\ \vdots \quad \ddots \quad \vdots \quad \vdots \\ \gamma_{a_1 a_{n-1}}^* \quad \cdots \quad \gamma_{a_{n-1} a_{n-1}}^* \quad \gamma_{b_n a_{n-1}}^* \\ \gamma_{a_1 a_n}^* \quad \cdots \quad \gamma_{a_{n-1} a_n}^* \quad \gamma_{b_n a_n}^* \end{array}.$$

4. Each pair of arrows $\delta: c \rightarrow a$, $\delta^*: a \rightarrow c$, $c \neq a$, $c \neq b$, is replaced with the ensemble

$$\Delta, \quad \Delta^* = c \begin{array}{c} a_1 \quad \cdots \quad a_{n-1} \quad a_n \\ \hline \delta_{a_1 c}^* \quad \cdots \quad \delta_{a_{n-1} c}^* \quad \delta_{a_n c}^* \end{array}.$$

The same is done for each pair of arrows $c \rightarrow b$ and $b \rightarrow c$, $c \neq a$, $c \neq b$.

The relations for the quiver $Q'_{\alpha,x}$ induced by the replacements are obtained from the relations for the quiver Q in the following way.

In every relation for the quiver Q , replace the morphism α with the morphism matrix


$$\begin{array}{c} a_1 \quad a_2 \quad \cdots \quad a_n \\ \begin{array}{|c|} \hline a_1 \quad x_1 \varepsilon_{a_1} \\ a_2 \quad \quad x_2 \varepsilon_{a_2} \quad 0 \\ \vdots \quad \quad 0 \quad \quad \ddots \\ a_n \quad \quad \quad \quad \quad \quad x_n \varepsilon_{a_n} \\ \hline \end{array} \end{array} \quad (1)$$

if $a = b$ and $\alpha = \alpha^*$. In the case where $\alpha: a \rightarrow b$ and $a \neq b$, replace α with the morphism matrix (1) if $x_i \neq 0$, $i \in \overline{1, n}$, and with the morphism matrix


$$\begin{array}{c} a_1 \quad a_2 \quad \cdots \quad a_{n-1} \quad a_n \\ \begin{array}{|c|} \hline a_1 \quad x_1 \varepsilon_{a_1} \\ a_2 \quad \quad x_2 \varepsilon_{a_2} \\ \vdots \quad \quad \quad \quad \quad \quad \ddots \\ a_{n-1} \quad \quad \quad \quad \quad \quad \quad \quad x_{n-1} \varepsilon_{a_{n-1}} \\ b_n \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad 0 \\ \hline \end{array} \end{array}$$

if $x_n = 0$. Also, replace the morphisms $\beta = \beta^*: a \rightarrow a$, $\tilde{\beta} = \tilde{\beta}^*: b \rightarrow b$, $\gamma: a \rightarrow b$, $\gamma^*: b \rightarrow a$, $\delta: c \rightarrow a$, $\delta^*: a \rightarrow c$ ($c \neq a$, $c \neq b$), $\tilde{\delta}: d \rightarrow b$, $\tilde{\delta}^*: b \rightarrow d$ ($d \neq a$, $d \neq b$) with the morphism matrices of the corresponding ensembles performing all the operations by formally multiplying and adding the matrices and equating the corresponding elements in the obtained identities. Naturally, one relation gives rise to several relations, in general.

Example 1. Let, for example, Q be the quiver




with the relation $\alpha^3 - 6\alpha^2 + 11\alpha - 6\varepsilon_a \equiv (\alpha - \varepsilon_a)(\alpha - 2\varepsilon_a)(\alpha - 3\varepsilon_a) = 0$, $x = (1, 2, 0)$. Then $Q'_{\alpha,x}$ has the form



and the relations are

$$\begin{array}{c} a_1 \ a_2 \ a_3 \\ a_1 \begin{bmatrix} 0 & & \\ & \varepsilon_{a_2} & \\ & & -\varepsilon_{a_3} \end{bmatrix} \cdot a_1 \begin{bmatrix} -\varepsilon_{a_1} & & \\ & 0 & \\ & & -2\varepsilon_{a_3} \end{bmatrix} \cdot a_1 \begin{bmatrix} -2\varepsilon_{a_1} & & \\ & -2\varepsilon_{a_2} & \\ & & -3\varepsilon_{a_3} \end{bmatrix} = 0. \\ a_2 \\ a_n \end{array}$$

Hence, $\varepsilon_{a_3} = 0$ and, consequently, a_3 is the zero object. Finally, $Q'_{\alpha,x}$ is the *-quiver



with no additional relations.

Remark 1. In a standard way, a not self-adjoint loop $\gamma: a \rightarrow a$ and its adjoint γ^* can be replaced with two self-adjoint ones by choosing another system of generators in the *-category $\mathcal{K}(Q)$, $\alpha = \frac{\gamma+\gamma^*}{2}$, $\beta = \frac{\gamma-\gamma^*}{2i}$. So, it is always possible to assume that the quiver Q has an arrow α with respect to which one can construct the derived quiver $Q'_{\alpha,x}$.

Theorem 1. For a quiver Q , let $Q'_{\alpha,x}$ be the derived quiver with respect to the arrow α at the point $x = (x_1, x_2, \dots, x_n)$. Then $\mathcal{K}(Q) \succ \mathcal{K}(Q'_{\alpha,x})$.

Proof. Let $\mathcal{M}(Q'_{\alpha,x})$ be the category of matrices over the *-category $\mathcal{K}(Q'_{\alpha,x})$. We will prove the claim in the case where $\alpha: a \rightarrow a$ is a self-adjoint arrow. If $\alpha: a \rightarrow b$, $a \neq b$, the proof is similar. Let us construct the functor $\Psi: \mathcal{K}(Q) \rightarrow \mathcal{M}(Q'_{\alpha,x})$ as follows. Set

$$\Psi(\alpha) = \begin{array}{c} a_1 \ a_2 \ \dots \ a_n \\ a_1 \begin{bmatrix} x_1 \varepsilon_{a_1} & & & \\ & x_2 \varepsilon_{a_2} & & \\ & & \ddots & \\ & & & x_n \varepsilon_{a_n} \end{bmatrix} \\ a_2 \\ \vdots \\ a_n \end{array}$$

$\Psi(\beta) = \beta$ if the arrow is not changed when constructing $Q'_{\alpha,x}$, and set $\Psi(\beta)$ to be the morphism matrix of the ensemble that replaces β , otherwise. Relations for $Q'_{\alpha,x}$ were defined so that the constructed mapping agreed with the relations for Q and $Q'_{\alpha,x}$ and could be extended to a functor from the category $\mathcal{K}(Q)$ into the category $\mathcal{K}(Q'_{\alpha,x})$.

Let us define the functor

$$\mathbf{E}_\Psi: \text{Rep}(Q'_{\alpha,x}, \mathcal{H}) \rightarrow \text{Rep}(Q, \mathcal{H})$$

as follows. If $\pi \in \text{Ob Rep}(Q'_{\alpha,x}, \mathcal{H})$, then π is extended to a representation Π of the category $\mathcal{M}(Q'_{\alpha,x})$ on \mathcal{H} , and we set $\mathbf{F}_\Psi = \Pi \circ \Psi$.

Let $\mathcal{C} = (\mathcal{C}_a)_{a \in Q'_v}$ be a morphism of representations in $\text{Rep}(Q'_{\alpha,x}, \mathcal{H})$, $\mathcal{C}_a: \pi(a) \rightarrow \pi_1(a)$. If $\bar{a} = (a_1, a_2, \dots, a_k) \in \text{Ob } \mathcal{M}(Q'_{\alpha,x})$, then set

$$\begin{aligned} (\mathbf{F}_\Psi(\mathcal{C}))_{\bar{a}} &= \text{diag}(\mathcal{C}_{a_1}, \mathcal{C}_{a_2}, \dots, \mathcal{C}_{a_k}) \\ &\in \text{Hom}_{\mathcal{M}(Q'_{\alpha,x})} \left((\pi(a_1), \pi(a_2), \dots, \pi(a_k)), (\pi_1(a_1), \pi_1(a_2), \dots, \pi_1(a_k)) \right). \end{aligned}$$

It is easy to check that $\mathbf{F}_\Psi(\mathcal{C})$ is a morphism from the representation $\mathbf{F}_\Psi(\pi)$ into the representation $\mathbf{F}_\Psi(\pi_1)$ (in the category $\text{Rep}(Q, \mathcal{H})$) and \mathbf{F}_Ψ is a functor. It is easy to check as well that, being univalent by the construction, the functor \mathbf{F}_Ψ is also full. Hence, \mathbf{F}_Ψ is an equivalence functor from the category $\text{Rep}(Q'_{\alpha,x}, \mathcal{H})$ into some full subcategory of the category $\text{Rep}(Q, \mathcal{H})$ and, by the definition of the majorization relation, we have $\mathcal{K}(Q) \succ \mathcal{K}(Q'_{\alpha,x})$. \square

It is sometimes useful to describe the repeated construction of the derived quiver with respect to arrows in an ensemble \mathcal{B} or a subquiver of a quiver Q as a single construction of the derived quiver $Q'_{\mathcal{B},x}$ with respect to the ensemble \mathcal{B} .

Example 2. Let Q be the ensemble \mathcal{A} and \mathcal{A}^* ,

$$\mathcal{A} = \begin{matrix} & a_1 & a_2 \\ b_1 & \boxed{\alpha_{11} \ \alpha_{12}} & \\ b_2 & \boxed{\alpha_{21} \ \alpha_{22}} & \end{matrix}, \quad \mathcal{A}^* = \begin{matrix} & b_1 & b_2 \\ a_1 & \boxed{\alpha_{11}^* \ \alpha_{21}^*} & \\ a_2 & \boxed{\alpha_{12}^* \ \alpha_{22}^*} & \end{matrix}$$

with the relation

$$\mathcal{A}^* \mathcal{A} = \begin{matrix} & a_1 & a_2 \\ a_1 & \boxed{\varepsilon_{a_1} \ 0} & \\ a_2 & \boxed{0 \ \varepsilon_{a_2}} & \end{matrix}.$$

Consider a subensemble $\mathcal{B}, \mathcal{B}^*$ in Q , where

$$\mathcal{B}^* = a_2 \begin{matrix} b_1 & b_2 \\ \boxed{\alpha_{12}^* \ \alpha_{22}^*} \end{matrix}.$$

Let $Q^{(1)} = Q'_{\mathcal{B},x}$ be the derived quiver for the quiver Q with respect to the subensemble \mathcal{B} at the point x that can be conveniently written as the matrix

$$x = \begin{array}{c} \begin{array}{c} a_2 \\ \begin{array}{|c|} \hline x_1 \\ \hline \end{array} \\ b_1 \end{array} \begin{array}{|c|} \hline \begin{array}{cc} x_2 & 0 \\ 0 & \ddots \\ & x_m \end{array} \\ \hline \end{array}, \\ \begin{array}{c} \begin{array}{|c|} \hline y_1 \\ \hline \end{array} \\ b_2 \end{array} \begin{array}{|c|} \hline \begin{array}{cc} y_2 & 0 \\ 0 & \ddots \\ & y_m \end{array} \\ \hline \end{array} \end{array}$$

where $x_i > 0, y_i > 0, x_i \neq x_j, y_i \neq y_j$ for $i \neq j, x_i^2 + y_i^2 = 1$. We wrote the point x as a matrix as to stress that, when constructing $Q'_{\mathcal{B},x}$, the ensemble $\mathcal{B}, \mathcal{B}^*$ is replaced with the matrix

$$\begin{array}{c} \begin{array}{c} a_2^{(1)} \\ a_2^{(2)} \\ \vdots \\ a_2^{(m)} \end{array} \begin{array}{|c|} \hline \begin{array}{ccc} x_1 \varepsilon_{a_2^{(1)}} & & \\ & x_2 \varepsilon_{a_2^{(2)}} & 0 \\ & 0 & \ddots \\ & & x_m \varepsilon_{a_2^{(m)}} \end{array} \\ \hline \end{array} \\ \begin{array}{c} a_2^{(1)} \\ a_2^{(2)} \\ \vdots \\ a_2^{(m)} \end{array} \begin{array}{|c|} \hline \begin{array}{ccc} y_1 \varepsilon_{a_2^{(1)}} & & \\ & y_2 \varepsilon_{a_2^{(2)}} & 0 \\ & 0 & \ddots \\ & & y_m \varepsilon_{a_2^{(m)}} \end{array} \\ \hline \end{array} \end{array}$$

and its adjoint.

The subensemble

$$\tilde{\mathcal{A}}^* = a_1 \begin{array}{|c|} \hline \begin{array}{cc} b_1 & b_2 \\ \alpha_{11}^* & \alpha_{12}^* \end{array} \\ \hline \end{array}, \quad \tilde{\mathcal{A}}$$

in the ensemble \mathcal{A}^* , \mathcal{A} will be replaced with

$$\mathcal{A}^{(1)*} = a_1 \begin{array}{c} a_2^{(1)} \ a_2^{(2)} \ \dots \ a_2^{(m)} \\ \hline \alpha_{11}^{(1)*} \ \alpha_{11}^{(2)*} \ \dots \ \alpha_{11}^{(m)*} \end{array} \begin{array}{c} a_2^{(1)} \ a_2^{(2)} \ \dots \ a_2^{(m)} \\ \hline \alpha_{11}^{(1)*} \ \alpha_{11}^{(2)*} \ \dots \ \alpha_{11}^{(m)*} \end{array}, \quad \mathcal{A}^{(1)}.$$

The relations in the quiver Q imply that

$$x_i \alpha_{11}^{(i)} + y_i \alpha_{21}^{(i)} = 0, \quad \sum_{i=1}^m (\alpha_{11}^{(1)*} \alpha_{11}^{(i)} + \alpha_{21}^{(i)*} \alpha_{21}^{(i)}) = \varepsilon_{a_1}$$

or, since $\alpha_{21}^{(i)} = -\frac{x_i}{y_i} \alpha_{11}^{(i)}$, we get

$$\sum_{i=1}^m \frac{\alpha_{11}^{(i)*} \alpha_{11}^{(i)}}{y_i^2} = \varepsilon_{a_1}.$$

If the system of generators in the category $\mathcal{K}(Q^{(1)})$ is changed by replacing the arrows $\alpha_{11}^{(i)}$ with $\beta_i = \frac{\alpha_{11}^{(i)}}{y_i}$, then we get that the category $\mathcal{K}(Q^{(1)})$ is generated by the ensemble

$$\mathcal{A}^{(2)*} = a_1 \begin{array}{c} a_2^{(1)} \ a_2^{(2)} \ \dots \ a_2^{(m)} \\ \hline \beta_{11}^{(1)*} \ \beta_{11}^{(2)*} \ \dots \ \beta_{11}^{(m)*} \end{array}, \quad \mathcal{A}^{(2)}$$

and the relation $\mathcal{A}^{(2)*} \cdot \mathcal{A}^{(2)} = \varepsilon_{a_1}$.

4. ENSEMBLES WITH AN ORTHOGONALITY CONDITION

We will call an ensemble

$$\mathcal{A} = \begin{array}{c} b_1 \\ b_2 \\ \vdots \\ b_m \end{array} \begin{array}{c} a_1 \ a_2 \ \dots \ a_n \\ \hline \alpha_{11} \ \alpha_{12} \ \dots \ \alpha_{1n} \\ \alpha_{21} \ \alpha_{22} \ \dots \ \alpha_{2n} \\ \vdots \ \quad \quad \quad \ddots \ \quad \quad \quad \vdots \\ \alpha_{m1} \ \alpha_{m2} \ \dots \ \alpha_{mn} \end{array}, \quad \mathcal{A}^*$$

with the relation

$$\mathcal{A}^* \mathcal{A} = \begin{matrix} & a_1 & a_2 & \cdots & a_n \\ \begin{matrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{matrix} & \boxed{\begin{matrix} \varepsilon_{a_1} & & & \\ & \varepsilon_{a_2} & & 0 \\ & & \ddots & \\ & 0 & & \varepsilon_{a_n} \end{matrix}} & , & \alpha_{ij} : a_j \rightarrow b_i, \end{matrix}$$

an ensemble of dimension $m \times n$ with the orthogonality condition

$$\alpha_{1i}^* \alpha_{1j} + \alpha_{2i}^* \alpha_{2j} + \cdots + \alpha_{mi}^* \alpha_{mj} = \begin{cases} 0, & \text{if } i \neq j, \\ \varepsilon_{a_i}, & \text{if } i = j \end{cases}$$

in the vertices a_1, a_2, \dots, a_n and denote it by $Q_{m \times n^\perp}$.

Theorem 2. *The following relations take place for the categories:*

1. $\mathcal{K}(Q_{k \times l^\perp}) \succ \mathcal{K}(Q_{m \times n^\perp})$ for $k \geq 2, l \geq 2$, and any $m, n \in \mathbb{N}$;
2. $\mathcal{K}(Q_{k \times 1^\perp}) \succ \mathcal{K}(Q_{m \times n^\perp})$ for $k \geq 3$ and any $m, n \in \mathbb{N}$.

Proof. Since the majorization relation is transitive and $\mathcal{K}(Q_{m_1 \times n_1^\perp}) \succ \mathcal{K}(Q_{m_2 \times n_2^\perp})$ for $m_1 \geq m_2$ and $n_1 \geq n_2$, it remains to prove that

- a) $\mathcal{K}(Q_{2 \times 2^\perp}) \succ \mathcal{K}(Q_{m \times n^\perp})$;
- b) $\mathcal{K}(Q_{(m+1) \times 1^\perp}) \succ \mathcal{K}(Q_{m \times n^\perp})$;
- c) $\mathcal{K}(Q_{3 \times 1^\perp}) \succ \mathcal{K}(Q_{2 \times 2^\perp})$.

Ad a) Follows from Example 2 and Theorem 1.

Ad b) Suppose we have an ensemble $\mathcal{K}(Q_{(m+1) \times 1^\perp})$ given by the matrix

$$\mathcal{A}^* = a \begin{matrix} & b_1 & b_2 & \cdots & b_{m+1} \\ \boxed{\alpha_1^* \alpha_2^* \cdots \alpha_{m+1}^*} & & & & \end{matrix}, \quad \mathcal{A}$$

and the relation $\mathcal{A}^* \mathcal{A} = \varepsilon_a$. Construct the derived quiver $Q' = Q'_{\alpha_{m+1}, x}$, where $x = (x_1, x_2, \dots, x_n)$, $x_i \neq x_j$ for $i \neq j$, $0 < x_i < 1$. As a result we get the ensemble

$$\mathcal{B} = \begin{matrix} & a_1 & \cdots & a_n \\ \begin{matrix} b_1 \\ \vdots \\ b_m \end{matrix} & \boxed{\begin{matrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & \ddots & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{matrix}} & , & \mathcal{B}^* \end{matrix}$$

with the relation

$$\mathcal{B}^* \mathcal{B} = \begin{array}{c} a_1 \\ \vdots \\ a_n \end{array} \begin{array}{c} a_1 \quad \cdots \quad a_n \\ \boxed{\begin{array}{ccc} (1-x_1^2)\varepsilon_{a_1} & & \\ & \ddots & 0 \\ & & (1-x_n^2)\varepsilon_{a_n} \end{array}} \end{array}.$$

By replacing, in the category $\mathcal{K}(Q')$, the generators α_{ij} with $\beta_{ij} = \alpha_{ij}/\sqrt{1-x_j^2}$, we get the ensemble $Q_{m \times n^\perp}$. Consequently, by Theorem 1, we see that $\mathcal{K}(Q_{(m+1) \times 1^\perp}) \succ \succ \mathcal{K}(Q_{m \times n^\perp})$.

Ad c) Let $Q = Q_{3 \times 1^\perp}$. As in b), construct the derived quiver $Q'_{\alpha_3, x}$, where $x = (x_1, x_2)$, $x_1 \neq x_2$, $0 < x_i < 1$. We see that $\mathcal{K}(Q') = \mathcal{K}(Q_{2 \times 2^\perp})$ and, consequently, $\mathcal{K}(Q_{3 \times 1^\perp}) \succ \succ \mathcal{K}(Q_{2 \times 2^\perp})$ by Theorem 1. \square

We also have the following theorem.

Theorem 3. *The category $\mathcal{K}(Q_{k \times l^\perp})$ is $*$ -wild if and only if $k \geq 2$ and $l \geq 2$, or $k \geq 3$.*

Proof. The proof is actually given in [1] when proving Theorem 4.

Let $\mathcal{F}_2 = \mathbb{C} \langle u_1, u_2 \mid u_i u_i^* = u_i^* u_i = e \rangle$. Let us show that $\mathcal{K}(Q_{3 \times 1^\perp}) \succ C^*(\mathcal{F}_2)$. Suppose $\mathcal{K}(Q_{3 \times 1^\perp})$ is given by the ensemble \mathcal{A}^* , \mathcal{A} with

$$\mathcal{A}^* = a \begin{array}{c} b_1 \quad b_2 \quad b_3 \\ \boxed{\alpha_1^* \quad \alpha_2^* \quad \alpha_3^*} \end{array}.$$

Set

$$\Psi(\alpha_1) = \begin{array}{c} b_1^{(1)} \\ b_1^{(2)} \\ b_1^{(3)} \\ b_1^{(4)} \end{array} \begin{array}{c} a_1 \quad a_2 \quad a_3 \quad a_4 \quad a_5 \\ \boxed{\begin{array}{ccccc} e & 0 & 0 & 0 & 0 \\ 0 & e & 0 & 0 & 0 \\ 0 & 0 & 2e & 0 & 0 \\ 0 & 0 & 0 & 3e & 0 \end{array}} \end{array} \cdot \frac{1}{N}, \quad \Psi(\alpha_2) = \begin{array}{c} b_2^{(1)} \\ b_2^{(2)} \\ b_2^{(3)} \end{array} \begin{array}{c} a_1 \quad a_2 \quad a_3 \quad a_4 \quad a_5 \\ \boxed{\begin{array}{ccccc} e & 0 & e & e & e \\ 0 & 2e & e & u_1 & 0 \\ 0 & 0 & e & 0 & u_2 \end{array}} \end{array} \cdot \frac{1}{N},$$

$$\Psi(\alpha_3) = \sqrt{I - \Psi(\alpha_1)^* \Psi(\alpha_1) - \Psi(\alpha_2)^* \Psi(\alpha_2)},$$

where $I = \text{diag} \{e, e, e, e, e\}$ and N is chosen so that $\|\Psi(\alpha_1)^* \Psi(\alpha_1) - \Psi(\alpha_2)^* \Psi(\alpha_2)\| < < 1$ in the matrix algebra $\mathcal{M}_5(C^*(\mathcal{F}_2))$. It can be directly checked that the functor $\mathbf{F}_\Psi: \text{Rep}(C^*(\mathcal{F}_2)) \rightarrow \text{Rep}(Q_{3 \times 1^\perp})$ is univalent and full. Finally, the proof of the theorem follows from Theorem 2. \square

Acknowledgements

This work was supported by the Ukrainian State Fund for Fundamental Studies, No. 01.07/071.

REFERENCES

- [1] Kruglyak S. A., Samoilenko Yu. S.: *On structure theorems for a family of idempotents*. Ukrainskii Matematicheskii Zhurnal **50**(4), (1998) (Russian).
- [2] Kruglyak S. A., Samoilenko Yu. S.: *On the complexity of description of representations of *-algebras generated by idempotents*. [in:] Proc. of the American Mathematical Society, vol. 128, 1655–1664, AMS, 2000.
- [3] Kruglyak S. A.: *A majorization relation for *-categories and *-wild categories*. [in:] Proc. of the Fifth Intern. Conference “Symmetry in Nonlinear Math. Physics”, 2004.
- [4] Roiter A. V.: *Boxes with an involution*. [in:] Representations and quadratic forms, Kiev, 1979, 124–126 (Russian).
- [5] Kruglyak S. A.: *Representations of free involutive quivers*. [in:] Representations and quadratic forms, Kiev, 1979, 149–151 (Russian).
- [6] Kruglyak S. A., Samoilenko Yu. S.: *On unitary equivalence of collections of self-adjoint operators*. Funct. Anal. Appl. 1980, 84–85, (Russian).

O. V. Bagro, S. A. Kruglyak

Institute of Mathematics of the National Academy of Sciences of Ukraine,
vul. Tereshchinkivs'ka, 3, Kyiv, 01601, Ukraine

Received: January 5, 2004.