

You Lu, Xinmin Hou, Jun-Ming Xu

A NOTE ON THE p -DOMINATION NUMBER OF TREES

Abstract. Let p be a positive integer and $G = (V(G), E(G))$ a graph. A p -dominating set of G is a subset S of $V(G)$ such that every vertex not in S is dominated by at least p vertices in S . The p -domination number $\gamma_p(G)$ is the minimum cardinality among the p -dominating sets of G . Let T be a tree with order $n \geq 2$ and $p \geq 2$ a positive integer. A vertex of $V(T)$ is a p -leaf if it has degree at most $p - 1$, while a p -support vertex is a vertex of degree at least p adjacent to a p -leaf. In this note, we show that $\gamma_p(T) \geq (n + |L_p(T)| - |S_p(T)|)/2$, where $L_p(T)$ and $S_p(T)$ are the sets of p -leaves and p -support vertices of T , respectively. Moreover, we characterize all trees attaining this lower bound.

Keywords: p -domination number, trees.

Mathematics Subject Classification: 05C69.

1. INTRODUCTION

Let $G = (V(G), E(G))$ be a simple graph with vertex set $V(G)$ and edge set $E(G)$. The *Open neighborhood* of a vertex $v \in V(G)$, denoted by $N_G(v)$, is the set of adjacent vertices of v , and the *Closed neighborhood* $N_G[v] = N_G(v) \cup \{v\}$. Let $\deg_G(v) = |N_G(v)|$ denote the degree of v . The *maximum degree* $\Delta(G) = \max\{\deg_G(v) : v \in V(G)\}$. For $S \subseteq V(G)$, the subgraph induced by S is denoted by $G[S]$. For a pair of vertices $u, v \in V(G)$, the *distance* $d_G(u, v)$ between u and v is the length of the shortest uv -paths in G . The *diameter* of G is $d(G) = \max\{d_G(u, v) : u, v \in V(G)\}$.

Let T be a nontrivial tree and $p \geq 2$ a positive integer. A p -leaf of T is a vertex with degree at most $p - 1$, while a p -support vertex of T is a vertex of degree at least p adjacent to a p -leaf. We denote the sets of p -leaves and p -support vertices of T by $L_p(T)$ and $S_p(T)$, respectively. Notice that if $p = 2$ then the 2-leaves (resp. 2-support vertices) are the usual leaves (resp. support vertices) of T , while $L_2(T)$ (resp. $S_2(T)$) is the set of leaves (resp. support vertices) of T . A tree T is a double star if it contains exactly two vertices that are not leaves. A double star with two support vertices a and b is denoted by $S_{a,b}$.

For notation and graph theory terminology we follow [2, 5, 6]. For a vertex v in a rooted tree T , we let $C(v)$ and $D(v)$ denote the set of children and descendants, respectively, of v , and we define $D[v] = D(v) \cup \{v\}$.

In [4], Fink and Jacobson introduced the concept of p -domination. Let p be a positive integer. A subset S of $V(G)$ is a p -dominating set of G if for every vertex $v \in V(G) - S$, $|S \cap N_G(v)| \geq p$. The p -domination number $\gamma_p(G)$ is the minimum cardinality among the p -dominating sets of G . Any p -dominating set of G with cardinality $\gamma_p(G)$ will be called a γ_p -set of G . Note that the 1-domination number $\gamma_1(G)$ is the classical domination number $\gamma(G)$. For any $S, T \subseteq V(G)$, S p -dominates T in G if for every vertex $v \in T - S$, $|S \cap N_G(v)| \geq p$.

Some bounds of the p -domination number of a tree T are given in literature. Let T be a nontrivial tree of order $n \geq 3$ and with l leaves and s support vertices. Lemańska [7] showed that $\gamma(T) \geq (n + 2 - l)/2$. Chellali [3] proved that $\gamma_2(T) \geq (n + l - s)/2$ and this lower bound is sharp. In [1], Blidia et al. proved that, for $p \geq 2$, $\gamma_p(T) \leq (n + |L_p(T)|)/2$.

In this note, we give a lower bound of $\gamma_p(T)$ in terms of $n, |L_p(T)|, |S_p(T)|$, that is: Let T be a tree of order n . Then

$$\gamma_p(T) \geq (n + |L_p(T)| - |S_p(T)|)/2$$

for $p \geq 2$, which generalizes the lower bounds of Chellali [3]. Moreover, we characterize all trees attaining this lower bound.

Note that Fink and Jacobson [4] also provided a lower bound of $\gamma_p(T)$ in terms of the order n of a tree T and p , that is: $\gamma_p(T) \geq \frac{(p-1)n+1}{p}$. And recently Volkmann [8] characterized the family of trees with $\gamma_p(T) = \lceil \frac{(p-1)n+1}{p} \rceil$. Now we show that our bound is better than Fink and Jacobson's in some cases. Let $X_p(T)$ denote the set of vertices with degree at least p in T . Let $l_p = |L_p(T)|$ and $x_p = |X_p(T)|$. Then $\frac{\sum_{v \in L_p(T)} \text{deg}_T(v)}{l_p} \geq 1$ and $\frac{\sum_{v \in X_p(T)} \text{deg}_T(v)}{x_p} \geq p$. Let T be a tree satisfying the following conditions:

$$\frac{\sum_{v \in L_p(T)} \text{deg}_T(v)}{l_p} \geq 1 + \alpha \text{ and } \frac{\sum_{v \in X_p(T)} \text{deg}_T(v)}{x_p} \geq p + \beta, \tag{1.1}$$

where α and β are any two nonnegative constants satisfying $(p - 1)\alpha + \beta = 1$. Hence $2n - 2 = \sum_{v \in L_p(T)} \text{deg}_T(v) + \sum_{v \in X_p(T)} \text{deg}_T(v) \geq (1 + \alpha)l_p + (p + \beta)x_p = (1 + \alpha)n + p(1 - \alpha)x_p$, the second equality holds since $l_p + x_p = n$ and $(p - 1)\alpha + \beta = 1$. So $x_p \leq \frac{n}{p} - \frac{2}{p(1-\alpha)}$. Therefore,

$$\frac{n + |L_p(T)| - |S_p(T)|}{2} \geq \frac{n + l_p - x_p}{2} = n - x_p \geq n - \frac{n}{p} + \frac{2}{p(1-\alpha)} > \frac{(p-1)n+1}{p}.$$

This implies that our bound is better than the bound given by Fink and Jacobson for the trees satisfying condition (1.1). [Such trees satisfying condition (1.1) exist. For example, let T be a tree of order $n \geq 2$ and $cor_p(T)$ a tree obtained from T by adding p pendant edges to each vertex of T . Then $L_p(cor_p(T)) = V(cor_p(T)) - V(T)$ and

$X_p(\text{cor}_p(T)) = V(T)$. Let $T' = \text{cor}_p(T)$. Then we have $\frac{\sum_{v \in L_p(T')} \text{deg}_{T'}(v)}{l_p} = 1 = 1 + 0$ and $\frac{\sum_{v \in X_p(T')} \text{deg}_{T'}(v)}{x_p} = p + \frac{1}{n} \sum_{v \in V(T)} \text{deg}_T(v) = p + \frac{2n-2}{n} \geq p + 1.$

2. MAIN RESULTS

The following result is straightforward and can be found in [1].

Lemma 2.1. ([1]) *Every p -dominating set of a graph G contains any vertex of degree at most $p - 1$.*

A vertex is a central vertex of a star $K_{1,t}$ ($t \geq 1$) if either $t \geq 2$ and it is the support vertex or $t = 1$ and it is one of the two leaves. For convenience, we call an isolated vertex a star, denoted by $K_{1,0}$, and the only vertex is called the central vertex.

We define the family \mathcal{T}_p as:

$\mathcal{T}_p = \{T : T \text{ is obtained from a sequence } T_1, T_2, \dots, T_k \text{ (} k \geq 1 \text{) of trees, where } T_1 = K_{1,t} \text{ (} t \geq p \text{), } T = T_k \text{, and, if } k \geq 2 \text{, } T_{i+1} \text{ (} 1 \leq i \leq k - 1 \text{) is obtained from } T_i \text{ by using Operation } \mathcal{O}_j \text{ (} j = 1, 2 \text{ or } 3 \text{) listed below.}\}$

- **Operation \mathcal{O}_1** : Attach a copy of $K_{1,t}$ ($0 \leq t \leq p - 2$) by joining the central vertex to a p -support vertex of T_i or to a vertex of degree at most $p - 2$ in T_i .
- **Operation \mathcal{O}_2** : Attach a copy of $K_{1,t}$ ($t \geq p$) by joining the central vertex to a p -support vertex of T_i .
- **Operation \mathcal{O}_3** : Attach a copy of $K_{1,t}$ ($t \geq p - 1$) by joining the central vertex to a vertex of degree at most $p - 2$ in T_i .

From the way in which a tree $T \in \mathcal{T}_p$ is constructed we make the following lemma.

Lemma 2.2. *Let T be a tree in the family \mathcal{T}_p with $p \geq 2$. Then, $V(T) = L_p(T) \cup S_p(T)$ and each vertex of $S_p(T)$ is adjacent to at least p p -leaves in T .*

From Lemma 2.1 and Lemma 2.2, it is straightforward to obtain the follow result.

Lemma 2.3. *For any positive integer $p \geq 2$, if $T \in \mathcal{T}_p$, then $L_p(T)$ is the unique γ_p -set of T , and*

$$\gamma_p(T) = |L_p(T)| = (|V(T)| + |L_p(T)| - |S_p(T)|)/2.$$

Theorem 2.4. *Let T be a tree with order $n \geq 2$ and $p \geq 2$ a positive integer. Then*

$$\gamma_p(T) \geq (n + |L_p(T)| - |S_p(T)|)/2$$

with equality if and only if either $\Delta(T) \leq p - 1$ or $T \in \mathcal{T}_p$.

Proof. Let $l_p = |L_p(T)|$ and $s_p = |S_p(T)|$. We proceed by induction on the order n . If $n = 2$, then $T = P_2$, and so $\Delta(T) \leq p - 1$. The result holds. This establishes the base case. Assume that the result is true for every tree T' with order $2 \leq |V(T')| = n' < n$ and let T be a tree of order n .

If $\Delta(T) \leq p-1$, then $l_p = n$, $s_p = 0$ and by Lemma 2.1, $\gamma_p(T) = n = (n+l_p-s_p)/2$. The result follows. Assume now that $\Delta(T) \geq p$, then $d(T) \geq 2$. If $d(T) = 2$, then $T = K_{1,t}$ ($t \geq p$) belongs to \mathcal{T}_p . By Lemma 2.3, $\gamma_p(T) = (n+l_p-s_p)/2$. If $d(T) = 3$, then T is a double star $S_{a,b}$ with $\deg_T(a) \geq p$ or $\deg_T(b) \geq p$. Without loss of generality, assume $\deg_T(a) \geq p$. If $2 \leq \deg_T(b) \leq p-1$, then T can be obtained recursively from $K_{1,t}$ ($t = \deg_T(a) \geq p$) by attaching $\deg_T(b) - 1$ vertices to one leaf of $K_{1,t}$. Hence T is obtained recursively from $K_{1,t}$ by using $\deg_T(b) - 1$ Operations \mathcal{O}_1 . Hence $T \in \mathcal{T}_p$ and, by Lemma 2.3, $\gamma_p(T) = (n+l_p-s_p)/2$. If $\deg_T(b) = p$, then, to p -dominate b , one of a, b must be contained in a p -dominating set of T . Hence $\gamma_p(T) \geq l_p + 1 = (n+l_p-s_p)/2 + 1 > (n+l_p-s_p)/2$. If $\deg_T(b) > p$ and $\deg_T(a) = p$, then, to p -dominate a , one of a, b must be contained in a p -dominating set of T . Hence $\gamma_p(T) \geq l_p + 1 = (n+l_p-s_p)/2 + 1 > (n+l_p-s_p)/2$. If $\deg_T(b) > p$ and $\deg_T(a) > p$, then T can be seen as a tree constructed from a star $K_{1,t}$ ($t = \deg_T(a) - 1 \geq p$) by using Operation \mathcal{O}_2 by attaching a star $K_{1,m}$ ($m = \deg_T(b) - 1 \geq p$) to vertex a . Hence $T \in \mathcal{T}_p$. By Lemma 2.3, $\gamma_p(T) = (n+l_p-s_p)/2$. Therefore, in the following we assume that T is a tree with $d(T) \geq 4$ and $\Delta(T) \geq p$.

We now root T at a vertex r of maximum eccentricity. Let $P = uvwxy \cdots r$ be a longest path such that $d(u, r) = d(T) \geq 4$ and $\deg_T(v)$ is as large as possible. Then, each vertex in $D(w) - C(w)$ has degree one.

Case 1. $\deg_T(v) \geq p$.

Let $T' = T - D[v]$ and S be a γ_p -set of T such that S contains the vertices of $D[v]$ as few as possible. Since every vertex of $D(v)$ has degree one, $D(v) \subseteq S$. Hence $v \notin S$ (otherwise, we can replace v by w and get a γ_p -set of T which contains fewer vertices of $D[v]$ than S , a contradiction). Thus $S \cap V(T')$ is a p -dominating set of T' , and so

$$\gamma_p(T) = |S| = |D(v)| + |S \cap V(T')| \geq |D(v)| + \gamma_p(T').$$

Subcase 1.1. $\deg_T(w) \neq p$.

Then $n' = n - |D[v]|$, $l'_p = l_p - |D(v)|$ and $s'_p = s_p - 1$. By the inductive hypothesis on T' ,

$$\gamma_p(T) \geq |D(v)| + \gamma_p(T') \geq |D(v)| + (n' + l'_p - s'_p)/2 = (n + l_p - s_p)/2.$$

Further, if $\gamma_p(T) = (n+l_p-s_p)/2$, then $\gamma_p(T') = (n'+l'_p-s'_p)/2$. By the inductive hypothesis on T' , $\Delta(T') \leq p-1$ or $T' \in \mathcal{T}_p$.

If $\Delta(T') \leq p-1$, then $\deg_T(w) = \deg_{T'}(w) + 1 \leq \Delta(T') + 1 \leq p$. Since $\deg_T(w) \neq p$, $\deg_T(w) \leq p-1$, and so v is the unique vertex with degree at least p in T . Hence T can be obtained recursively from a star $K_{1,t}$ ($t = \deg_T(v) \geq p$) by attaching $n-t-1$ isolated vertices. Hence T is obtained recursively from the star $K_{1,t}$ by using $n-t-1$ operations \mathcal{O}_1 , and so $T \in \mathcal{T}_p$.

If $T' \in \mathcal{T}_p$, then, by Lemma 2.3, $S \cap V(T') = L_p(T')$. If $\deg_T(w) \leq p-1$, then $\deg_{T'}(w) \leq p-2$. Thus T is obtained from T' by using Operation \mathcal{O}_3 by attaching the star $K_{1,t}$ ($= T[D[v]]$, $t = \deg_T(v) - 1 \geq p-1$) to w . Hence $T \in \mathcal{T}_p$. If $\deg_T(w) \geq p+1$ and $\deg_T(v) \geq p+1$, then $\deg_{T'}(w) \geq p$. Thus T is obtained from T' by using Operation \mathcal{O}_2 by attaching the star $K_{1,t}$ ($= T[D[v]]$, $t = \deg_T(v) - 1 \geq p$) to w . Hence $T \in \mathcal{T}_p$. If $\deg_T(w) \geq p+1$ and $\deg_T(v) = p$, we claim that the

equality doesn't hold in this case. Since $\deg_T(w) \geq p + 1$, then $\deg_{T'}(w) \geq p$. Hence, $w \notin L_p(T') = S \cap V(T')$. Since $v \notin S$ and w p -dominates v , $w \in S$, a contradiction.

Subcase 1.2. $\deg_T(w) = p$.

Then $n' = n - |D[v]|$ and $l'_p = l_p - |D(v)| + 1$. Now we count the number of the p -support vertices of T' . Note that a p -support vertex of T in $C(w) \setminus \{v\}$ is a p -support vertex of T' , too. Define δ_1 to be equal to 1, if $w \in S_p(T)$; and 0, otherwise. Define δ_2 to be equal to 1, if $\deg_T(x) \geq p$ and $x \notin S_p(T)$; and 0, otherwise. Then $s'_p = s_p - 1 - \delta_1 + \delta_2$. By the inductive hypothesis on T' ,

$$\begin{aligned} \gamma_p(T) &\geq |D(v)| + \gamma_p(T') \geq \\ &\geq |D(v)| + (n' + l'_p - s'_p)/2 = \\ &= (n + l_p - s_p)/2 + (\delta_1 + 1 - \delta_2)/2 \geq \\ &\geq (n + l_p - s_p)/2. \end{aligned}$$

We claim that the equality doesn't hold in this case. Suppose to the contrary that $\gamma_p(T) = (n + l_p - s_p)/2$. Then $\delta_1 + 1 = \delta_2$, and $\gamma_p(T') = (n' + l'_p - s'_p)/2$. By the definitions of δ_1 and δ_2 , $\delta_1 = 0$ and $\delta_2 = 1$. So $\deg_T(x) \geq p$ and $x \notin S_p(T)$. By inductive hypothesis on T' , $T' \in \mathcal{T}_p$. By Lemma 2.3, $L_p(T')$ is the unique p -dominating set of T' . Since $\deg_{T'}(w) = p - 1$, w is a p -leaf of T' . Hence x is a p -support vertex of T' . By Lemma 2.2, x be adjacent to at least p p -leaves of T' , and so x must be adjacent to at least $p - 1$ (≥ 1) p -leaves in T . Thus x is a p -support vertex of T since $\deg_T(x) \geq p$, which contradicts $x \notin S_p(T)$.

Case 2. $\deg_T(v) \leq p - 1$

By our choice of the path $P = uvwxy \dots r$, each vertex in $D(w) - C(w)$ has degree one, and for each vertex $a \in C(w)$, $\deg_T(a) \leq \deg_T(v) \leq p - 1$. Hence $D(w) \subseteq L_p(T)$.

Subcase 2.1. $\deg_T(w) \leq p - 1$ or $\deg_T(w) \geq p + 2$.

Let $T' = T - D[v]$. Let S be a γ_p -set of T . Then $n' = n - |D[v]|$, $l'_p = l_p - |D[v]|$ and $s'_p = s_p$. If $\deg_T(w) \leq p - 1$, then, by Lemma 2.1, $w \in S$. Hence $S \cap V(T')$ is a p -dominating set of T' . If $\deg_T(w) \geq p + 2$, then $C(w) \setminus \{v\} \subseteq D(w) \subseteq L_p(T) \subseteq S$. Hence $C(w) \setminus \{v\} \subseteq S \cap V(T')$. Since $|C(w)| - 1 = |\deg_T(w)| - 2 \geq p$, w is p -dominated by $S \cap V(T')$ and hence $S \cap V(T')$ is a p -dominating set of T' , too. By the induction on T' ,

$$\begin{aligned} \gamma_p(T) &= |S| = |D[v]| + |S \cap V(T')| \geq \\ &\geq |D[v]| + \gamma_p(T') \geq \\ &\geq |D[v]| + (n' + l'_p - s'_p)/2 = \\ &= (n + l_p - s_p)/2. \end{aligned}$$

Further if $\gamma_p(T) = (n + l_p - s_p)/2$, then $\gamma_p(T') = (n' + l'_p - s'_p)/2$. By the inductive hypothesis on T' , $\Delta(T') \leq p - 1$ or $T' \in \mathcal{T}_p$. We claim that the equality does not hold for $\Delta(T') \leq p - 1$. If not, then $\deg_T(w) = \deg_{T'}(w) + 1 \leq \Delta(T') + 1 \leq p$. Since $\deg_T(w) \leq p - 1$ or $\deg_T(w) \geq p + 2$, we have $\deg_T(w) \leq p - 1$. Thus $\Delta(T) \leq p - 1$, which contradicts the assumption that $\Delta(T) \geq p$.

Note that $\deg_{T'}(w) \leq p-2$ or $\deg_{T'}(w) \geq p+1$ ($w \in S_p(T')$ since $D(w) \subseteq L_p(T)$). Since $T' \in \mathcal{T}_p$, T can be constructed from T' by Operation \mathcal{O}_1 by attaching the star $K_{1,t}$ ($= T[D[v]]$, $t = \deg_T(v) - 1 \leq p-2$) to w . Hence, $T \in \mathcal{T}_p$.

Subcase 2.2. $\deg_T(w) = p$ and $\deg_T(x) \neq p$ or $\deg_T(w) = p+1$ and $\deg_T(x) \neq p$.

Let $T' = T - D[w]$, then $n' = n - |D[w]|$ and $l'_p = l_p - |D(w)|$ (since $D(w) \subseteq L_p(T)$ and $\deg_T(x) \neq p$). Since $\deg_T(x) \neq p$, $S_p(T') = S_p(T) \setminus \{w\}$ and $s'_p = s_p - 1$. Let S be a γ_p -set of T that contains the vertices of $D[w]$ as few as possible. Then $w \notin S$ (otherwise, we can replace w by x). Hence $S \cap V(T')$ is a p -dominating set of T' . By the induction on T' ,

$$\begin{aligned} \gamma_p(T) &= |S| = |D(w)| + |S \cap V(T')| \geq \\ &\geq |D(w)| + \gamma_p(T') \geq \\ &\geq |D(w)| + (n' + l'_p - s'_p)/2 = \\ &= (n + l_p - s_p)/2. \end{aligned}$$

Further if $\gamma_p(T) = (n + l_p - s_p)/2$, then $\gamma_p(T') = (n' + l'_p - s'_p)/2 = |S \cap V(T')|$. Hence $S \cap V(T')$ is a γ_p -set of T' . By the induction on T' , $\Delta(T') \leq p-1$ or $T' \in \mathcal{T}_p$. If $\Delta(T') \leq p-1$, then $\deg_T(w) = \deg_{T'}(w) + 1 \leq \Delta(T') + 1 \leq p$. Since $\deg_T(w) = p$ or $p+1$, $\deg_T(w) = p$. Thus w is a unique vertex of degree at least p in T . Hence T can be obtained recursively from $K_{1,t}$ ($t = \deg_T(w) = p$) by attaching $n-t-1$ isolated vertices. Hence T can be obtained recursively from $K_{1,t}$ by using $n-t-1$ operations \mathcal{O}_1 , and so $T \in \mathcal{T}_p$. Assume now that $T' \in \mathcal{T}_p$. By Lemma 2.3, $S \cap V(T') = L_p(T')$.

If $\deg_T(w) = p$ and $\deg_T(x) \geq p+1$, we claim that the equality doesn't hold in this case. Since $\deg_T(x) \geq p+1$, $\deg_{T'}(x) = \deg_T(x) - 1 \geq p$ and so $x \notin L_p(T') = S \cap V(T')$. Hence $x \notin S$. Note that $\deg_T(w) = p$ and $w \notin S$, to p -dominate w , $x \in S$, a contradiction.

If $\deg_T(w) = p+1$ and $\deg_T(x) \geq p+1$, then let $T'' = T[V(T) - (D(w) - C(w))]$. Note that $\deg_{T'}(x) = \deg_T(x) - 1 \geq p$, by Lemma 2.2, $x \in S_p(T')$. Thus T'' is obtained from T' by using Operation \mathcal{O}_2 by attaching the star $K_{1,t}$ ($= T[C(w) \cup \{w\}]$, $t = \deg_T(w) - 1 = p$) to x , and so $T'' \in \mathcal{T}_p$. By $2 \leq \deg_T(v) \leq p-1$, $p \geq 3$. Since $T[D(w) - C(w)]$ consists of $|D(w) - C(w)|$ isolated vertices, T is obtained recursively from T'' by attaching $|D(w) - C(w)|$ isolated vertices to some vertices of $C(w)$. Hence T is obtained recursively from T'' by using $|D(w) - C(w)|$ operations \mathcal{O}_1 , and so $T \in \mathcal{T}_p$.

If $\deg_T(x) \leq p-1$, then $p \geq 3$ and let $T'' = T[V(T) - (D(w) - C(w))]$. Since $\deg_{T'}(x) = \deg_T(x) - 1 \leq p-2$. T'' is obtained from T' by using Operation \mathcal{O}_3 by attaching the star $K_{1,t}$ ($= T[C(w) \cup \{w\}]$, $t = \deg_T(w) - 1 \geq p-1$) to x . So, $T'' \in \mathcal{T}_p$. Since $T[D(w) - C(w)]$ consists of $|D(w) - C(w)|$ isolated vertices, T is obtained recursively from T'' by attaching $|D(w) - C(w)|$ isolated vertices to some vertices of $C(w)$. Hence T is obtained recursively from T'' by using $|D(w) - C(w)|$ operations \mathcal{O}_1 , and so $T \in \mathcal{T}_p$.

Subcase 2.3. $\deg_T(w) = p+1$ and $\deg_T(x) = p$.

Let $T' = T - D[v]$. Note that $D[v] \subseteq L_p(T)$, we have $n' = n - |D[v]|$, $l'_p = l_p - |D[v]|$ and $s'_p = s_p$. Let S be a γ_p -set of T , then by Lemma 2.1, $D[v] \subseteq S$. If $w \in S$, then $S \cap V(T')$ is a p -dominating set of T' . Since $\deg_T(x) = p$ and S p -dominates w ,

$w \in S$ or $x \in S$. If $w \notin S$, then, to p -dominate x , $x \in S$ since $\deg_T(x) = p$. Hence $|(S \cap V(T')) \cap N_{T'}(w)| = |(C(w) - \{v\}) \cup \{x\}| = \deg_T(w) - 1 = p$. So, w is p -dominated by $S \cap V(T')$. Thus $S \cap V(T')$ is a p -dominating set of T' . By the induction on T' ,

$$\begin{aligned} \gamma_p(T) &= |S| = |D[v]| + |S \cap V(T')| \geq \\ &\geq |D[v]| + \gamma_p(T') \geq \\ &\geq |D[v]| + (n' + l'_p - s'_p)/2 = \\ &= (n + l_p - s_p)/2. \end{aligned}$$

We claim that the equality doesn't hold in this case. Suppose to the contrary that $\gamma_p(T) = (n + l_p - s_p)/2$. Then $\gamma_p(T') = (n' + l'_p - s'_p)/2 = |S \cap V(T')|$. Hence $S \cap V(T')$ is a γ_p -set of T' . Since $\deg_{T'}(w) = \deg_T(w) - 1 = p$, $\Delta(T') \geq p$. By the inductive hypothesis on T' , $T' \in \mathcal{T}_p$. By Lemma 2.3, $S \cap V(T') = L_p(T')$. Since $\deg_{T'}(w) = p$ and $\deg_{T'}(x) = \deg_T(x) = p$, by Lemma 2.2, $w \notin L_p(T')$ and $x \notin L_p(T')$. Hence $w, x \notin S$. But, to p -dominate w and x , at least one of w, x is contained by S , a contradiction.

Subcase 2.4. $\deg_T(w) = p$ and $\deg_T(x) = p$.

Since $d(T) \geq 4$, the father y of x in the rooted tree T exists. Let $C_p(x)$ be the set of children of x with degree at least p . Since $\deg_T(w) = p$, $w \in C_p(x)$. Let $C_p(x) = \{w_1, \dots, w_t\}$ ($t \geq 1$). Then, for $1 \leq i \leq t$, $D(w_i) \neq \emptyset$. By the choice of the path P , each vertex of $\cup_{i=1}^t D(w_i)$ has degree at most $p - 1$. So $\cup_{i=1}^t D(w_i) \subseteq L_p(T)$ and $C_p(x) \subseteq S_p(T)$. Define δ to be equal to 1, if $\deg_T(y) \geq p$ and $y \notin S_p(T)$; and 0, otherwise.

Let $T' = T - \cup_{i=1}^t D(w_i)$. Then we have $n' = n - |\cup_{i=1}^t D(w_i)|$, $l'_p = l_p - |\cup_{i=1}^t D(w_i)| + 1$ and $s'_p = s_p - t + \delta$. Let S be a γ_p -set of T that contains the vertices of $\cup_{i=1}^t D(w_i)$ as few as possible. Then $w_i \notin S$ for $i = 1, \dots, t$ (otherwise, we can replace w_i by x). Hence, to p -dominate w , x must be in S for $\deg_T(w) = p$. Thus $S \cap V(T')$ is a p -dominating set of T' . By the induction on T' ,

$$\begin{aligned} \gamma_p(T) &= |S| = |\cup_{i=1}^t D(w_i)| + |S \cap V(T')| \geq \\ &\geq \sum_{i=1}^t |D(w_i)| + \gamma_p(T') \geq \\ &\geq \sum_{i=1}^t |D(w_i)| + (n' + l'_p - s'_p)/2 = \\ &= (n + l_p - s_p)/2 + (1 - \delta)/2 \geq \\ &\geq (n + l_p - s_p)/2. \end{aligned}$$

We claim that the equality doesn't hold in this case. If $\gamma_p(T) = (n + l_p - s_p)/2$, then $\delta = 1$ and $\gamma_p(T') = (n' + l'_p - s'_p)/2 = |S \cap V(T')|$. Hence $\deg_{T'}(y) = \deg_T(y) \geq p$, $y \notin S_p(T)$ and $S \cap V(T')$ is a γ_p -set of T' . Thus $\Delta(T') \geq p$ and $y \in S_p(T')$. By the inductive hypothesis on T' , $T' \in \mathcal{T}_p$. By Lemma 2.3, $S \cap V(T') = L_p(T')$. Then $y \notin S \cap V(T')$. So $y \notin S$. Hence, to p -dominate y , there are at least p p -leaves in T' (and hence $p - 1 \geq 1$ p -leaves in T) that are adjacent to y . That is $y \in S_p(T)$, which contradicts to $y \notin S_p(T)$. \square

Acknowledgments

The work was supported by NNSF of China (No.10701068 and No.10671191).

REFERENCES

- [1] M. Blidia, M. Chellali, L. Volkmann, *Some bounds on the p -domination number in trees*, Discrete Math. 306 (2006), 2031–2037.
- [2] G. Chartrand, L. Lesniak, *Graphs & Digraphs*, 3rd ed., Chapman & Hall, London, 1996.
- [3] M. Chellali, *Bounds on the 2-domination number in cactus graphs*, Opuscula Math. 26 (2006), 5–11.
- [4] J.F. Fink, M.S. Jacobson, *n -Domination in graphs*, [in:] Y.Alavi, A.J.Schwenk (Eds.), Graph Theory with Applications to Algorithms and Computer Science, Wiley, New York, 1985, 283–300.
- [5] T.W. Haynes, S.T. Hedetniemi, P.J. Slater, *Fundamentals of domination in graphs*, New York, Marcel Deliker, 1998.
- [6] T.W. Haynes, S.T. Hedetniemi, P.J. Slater, *Domination in Graphs: Advanced Topics*, New York, Marcel Deliker, 1998.
- [7] M. Lemańska, *Lower bound on the domination number of a tree*, Discuss. Math. Graph Theory 24(2) (2004), 165–169.
- [8] L.Volkmann, *Some remarks on lower bounds on the p -domination number in trees*, J. Combin. Math. Combin. Comput. 61 (2007), 159–167.

You Lu

University of Science and Technology of China
Department of Mathematics
Hefei, Anhui, 230026, China

Xinmin Hou
xmhou@ustc.edu.cn

University of Science and Technology of China
Department of Mathematics
Hefei, Anhui, 230026, China

Jun-Ming Xu

University of Science and Technology of China
Department of Mathematics
Hefei, Anhui, 230026, China

Received: September 9, 2008.

Revised: March 4, 2009.

Accepted: March 10, 2009.