

## THE EQUALITY CASE IN SOME RECENT CONVEXITY INEQUALITIES

Gyula Maksa, Zsolt Páles

**Abstract.** In this paper, we investigate a functional equation related to some recently introduced and investigated convexity type inequalities.

**Keywords:** generalized convexity, affine functions, functional equations, extension theorem.

**Mathematics Subject Classification:** 39B52.

### 1. INTRODUCTION

In a recent paper [24] by Varošanec, a common generalization of convex and  $s$ -convex functions, Godunova-Levin functions, and  $\mathcal{P}$ -functions is introduced in the following way: Let  $I$  be a nonvoid subinterval of  $\mathbb{R}$  (the set of all real numbers),  $h : [0, 1] \rightarrow \mathbb{R}$  and  $f : I \rightarrow \mathbb{R}$  be real-valued functions satisfying the inequality

$$f(tx + (1 - t)y) \leq h(t)f(x) + h(1 - t)f(y) \quad (1.1)$$

for all  $x, y \in I$  and  $t \in ]0, 1[$ . An even more general notion, the so-called  $(T, h)$ -convexity, can be found in Hájzy [11]: Let  $X$  be a real or complex normed space,  $D \subset X$  be a nonempty convex set,  $\emptyset \neq T \subset [0, 1]$ , and  $h : T \rightarrow \mathbb{R}$  be a function. A function  $f : D \rightarrow \mathbb{R}$  is  $(T, h)$ -convex if (1.1) holds for all  $x, y \in D$  and  $t \in T$ . It is clear that this generalizes the concepts of convexity ( $h(t) = t$ ,  $t \in [0, 1]$ , [24], [21]), the Breckner-convexity ( $h(t) = t^s$ ,  $t \in ]0, 1[$ , for some  $s \in \mathbb{R}$ , [5], [6]), the Godunova-Levin functions ( $h(t) = t^{-1}$ ,  $t \in ]0, 1[$ , [10]), the  $\mathcal{P}$ -functions ( $h(t) = 1$ ,  $t \in [0, 1]$ , [18]), and the  $t$ -convexity ( $T = \{t, 1 - t\}$ ,  $h(t) = t$ ,  $h(1 - t) = 1 - t$ , where  $0 < t < 1$  is a fixed number, Kuhn [14]). For further related results see Burai-Hájzy [1, 2] and Burai-Hájzy-Juhász [3, 4].

In this note, we focus on the functional equation related to these convexity properties and give the solutions of the following problem. Let  $X$  be a real or complex topological vector space,  $D \subset X$  be a nonempty open set,  $T$  be a nonempty set,

and  $\alpha, \beta, a, b : T \rightarrow \mathbb{R}$  be given functions. The problem is to find all the solutions  $f : D \rightarrow \mathbb{R}$  of the functional equation

$$f(\alpha(t)x + \beta(t)y) = a(t)f(x) + b(t)f(y) \quad (x, y \in D, t \in T) \quad (1.2)$$

provided that  $D$  is  $(\alpha, \beta)$ -convex, that is,  $\alpha(t)x + \beta(t)y \in D$  whenever  $x, y \in D$  and  $t \in T$ . To avoid the trivialities and the unimportant cases, we suppose that there exists an element  $t_0 \in T$  such that

$$\alpha(t_0)\beta(t_0)a(t_0)b(t_0) \neq 0. \quad (1.3)$$

We refer to the solutions of (1.2) as  $(\alpha, \beta, a, b)$ -affine functions and the solutions  $f$  of the corresponding inequality

$$f(\alpha(t)x + \beta(t)y) \leq a(t)f(x) + b(t)f(y) \quad (x, y \in D, t \in T)$$

will be called  $(\alpha, \beta, a, b)$ -convex functions. Besides those convexity notions we listed above this is a generalization of  $(t, q)$ -convexity ( $T = \{t\}, \alpha(t) = t, \beta(t) = 1 - t, a(t) = q, b(t) = 1 - q$ , where  $t, q \in ]0, 1[$  are fixed numbers, Kuhn [15], Matkowski-Pycia [16]), and Orlicz  $s$ -convexity ( $T = [0, 1], \alpha(t) = t^s, \beta(t) = (1 - t)^s, a(t) = t, b(t) = 1 - t$  for all  $t \in T$  and for some  $s \geq 1$ , Orlicz [17], Hudzik-Maligranda [12]).

Our purpose is to describe the  $(\alpha, \beta, a, b)$ -affine functions. Throughout this paper  $X$  denotes a real or complex topological vector space. A function  $A : X \rightarrow \mathbb{R}$  is called additive if it satisfies the Cauchy functional equation

$$A(x + y) = A(x) + A(y) \quad (x, y \in X).$$

Given a subfield  $S \subseteq \mathbb{R}$ , a function  $\varphi : S \rightarrow \mathbb{R}$  is said to be a field-homomorphism if  $\varphi$  is additive and multiplicative on  $S$ , i.e.,

$$\varphi(s + t) = \varphi(s) + \varphi(t) \quad \text{and} \quad \varphi(st) = \varphi(s)\varphi(t) \quad (s, t \in S).$$

## 2. THE RESULTS

Our investigations are based on the following extension theorem which is an immediate consequence of Theorem 1 in Radó-Baker [19].

**Theorem 2.1.** *Let  $U$  be a nonempty, open, connected subset of  $X \times X$  and define the following sets*

$$\begin{aligned} U_0 &:= \{x + y \mid (x, y) \in U\}, \\ U_1 &:= \{x \mid \exists y \in X : (x, y) \in U\}, \quad \text{and} \\ U_2 &:= \{y \mid \exists x \in X : (x, y) \in U\}. \end{aligned}$$

*Suppose that the functions  $f_i : U_i \rightarrow \mathbb{R}$ , ( $i = 0, 1, 2$ ) satisfy the functional equation*

$$f_0(x + y) = f_1(x) + f_2(y)$$

for all  $(x, y) \in U$ . Then there exist a unique additive function  $A : X \rightarrow \mathbb{R}$  and a unique pair  $(c_1, c_2) \in \mathbb{R}^2$  such that

$$\begin{aligned} f_0(x) &= A(x) + c_1 + c_2 & (x \in U_0), \\ f_1(x) &= A(x) + c_1 & (x \in U_1), \text{ and} \\ f_2(x) &= A(x) + c_2 & (x \in U_2). \end{aligned}$$

An important consequence of the above theorem is the following result.

**Theorem 2.2.** Let  $\gamma, \delta, p, q \in \mathbb{R}$  and  $\emptyset \neq D \subset X$  be an open and connected set such that  $\gamma\delta pq \neq 0$  and  $\gamma x + \delta y \in D$  for all  $x, y \in D$ . Then the function  $f : D \rightarrow \mathbb{R}$  satisfies the functional equation

$$f(\gamma x + \delta y) = pf(x) + qf(y) \quad (x, y \in D) \tag{2.1}$$

if, and only if, there exist an additive function  $A : X \rightarrow \mathbb{R}$  and a constant  $c \in \mathbb{R}$  such that

$$\begin{aligned} A(\gamma x) &= pA(x) & (x \in X), \\ A(\delta x) &= qA(x) & (x \in X), \\ c(p + q - 1) &= 0, & \text{and} \\ f(x) &= A(x) + c & (x \in D). \end{aligned} \tag{2.2}$$

*Proof.* Equation (2.1) implies that

$$f(x + y) = pf\left(\frac{1}{\gamma}x\right) + qf\left(\frac{1}{\delta}y\right) \quad (x \in \gamma D, y \in \delta D).$$

Applying Theorem 2.1 for the open and connected set  $U := (\gamma D) \times (\delta D)$  and the triplet of functions

$$\begin{aligned} f_0(x) &:= f(x), \quad x \in \gamma D + \delta D \subset D, \\ f_1(x) &:= pf\left(\frac{1}{\gamma}x\right), \quad x \in \gamma D, \\ f_2(x) &:= qf\left(\frac{1}{\delta}x\right), \quad x \in \delta D, \end{aligned}$$

we obtain that

$$pf\left(\frac{1}{\gamma}x\right) = A_0(x) + c_0 \quad (x \in \gamma D)$$

with some additive function  $A_0 : X \rightarrow \mathbb{R}$  and  $c_0 \in \mathbb{R}$ . Thus

$$f(x) = \frac{1}{p}A_0(\gamma x) + \frac{c_0}{p} \quad (x \in D),$$

whence, with the definitions  $A(x) := \frac{1}{p}A_0(\gamma x)$ ,  $x \in X$  and  $c := \frac{c_0}{p}$ ,

$$f(x) = A(x) + c \quad (x \in D)$$

follows.

Obviously,  $A : X \rightarrow \mathbb{R}$  is additive. Replacing this form of  $f$  into (2.1), we find that

$$A(\gamma x) - pA(x) + A(\delta y) - qA(y) - c(p + q - 1) = 0 \quad (x, y \in D).$$

This shows that, for all fixed  $y \in D$ , the polynomial function

$$x \mapsto A(\gamma x) - pA(x) + A(\delta y) - qA(y) - c(p + q - 1) \quad (x \in X)$$

vanishes on  $D$ , therefore it vanishes everywhere on  $X$  (see Székelyhidi [23]). This implies the other equalities of (2.2), as well. The converse is straightforward.  $\square$

In the result below we investigate homogeneity properties of additive functions. Given an additive function  $A : \mathbb{R} \rightarrow \mathbb{R}$ , we introduce its set of homogeneity pairs  $H_A$  as follows:

$$H_A := \{(s, t) \in \mathbb{R}^2 \mid A(sx) = tA(x) \text{ for all } x \in \mathbb{R}\}.$$

**Theorem 2.3.** *Let  $A : \mathbb{R} \rightarrow \mathbb{R}$  be a nonzero additive function. Then there exist a subfield  $S_A \subseteq \mathbb{R}$  (called the homogeneity field of  $A$ ) and an injective field-homomorphism  $\varphi_A : S_A \rightarrow \mathbb{R}$  (called the homogeneity field-homomorphism of  $A$ ) such that  $H_A$  is equal to the graph of  $\varphi_A$ , i.e.,*

$$H_A = \{(s, \varphi_A(s)) \mid s \in S_A\}. \quad (2.3)$$

Conversely, for every subfield  $S \subseteq \mathbb{R}$  and injective field-homomorphism  $\varphi : S \rightarrow \mathbb{R}$ , there exists a nonzero additive function  $A : X \rightarrow \mathbb{R}$  such that  $S \subseteq S_A$  and  $\varphi_A|_S = \varphi$ .

*Proof.* Denote by  $S_A$  the domain of the relation  $H_A$ . We show that,  $H_A$  is in fact a function. Assume that  $(s, t_1), (s, t_2) \in H_A$ . Then, for all  $x \in X$ ,

$$(t_1 - t_2)A(x) = t_1A(x) - t_2A(x) = A(sx) - A(sx) = 0,$$

which, by the nontriviality of  $A$ , yields that  $t_1 = t_2$  proving that the relation  $H_A$  is a function. This means that there exists a function  $\varphi_A : S_A \rightarrow \mathbb{R}$  such that (2.3) holds. It remains to show that  $S_A$  is a subfield of  $\mathbb{R}$  and  $\varphi_A$  is an injective field-homomorphism.

To prove the injectivity, let  $(s_1, t), (s_2, t) \in H_A$ . Then, for all  $x \in X$ ,

$$A((s_1 - s_2)x) = A(s_1x) - A(s_2x) = tA(x) - tA(x) = 0,$$

which, by the nontriviality of  $A$ , yields that  $s_1 = s_2$ . By the injectivity,  $\varphi_A(s)$  is nonzero whenever  $s$  is different from zero.

Let  $s, t \in S$ . Then, using (2.3), for all  $x \in X$ , we get that

$$A((s - t)x) = A(sx) - A(tx) = \varphi_A(s)A(x) - \varphi_A(t)A(x) = (\varphi_A(s) - \varphi_A(t))A(x).$$

Hence,  $(s - t, \varphi_A(s) - \varphi_A(t)) \in H_A$ , which yields that  $s - t \in S$  and  $\varphi_A(s - t) = \varphi_A(s) - \varphi_A(t)$ . Thus  $S$  is a group with respect to the addition and  $\varphi_A$  is additive.

Similarly, for all  $s \in S, t \in S \setminus \{0\}$ , and  $x \in X$ , we obtain that

$$\varphi_A(t)A\left(\frac{s}{t}x\right) = A(sx) = \varphi_A(s)A(x).$$

Hence  $\left(\frac{s}{t}, \frac{\varphi_A(s)}{\varphi_A(t)}\right) \in H_A$ , which yields that  $\frac{s}{t} \in S$  and  $\varphi_A\left(\frac{s}{t}\right) = \frac{\varphi_A(s)}{\varphi_A(t)}$ . This proves that  $S$  is a semigroup under the multiplication whose nonzero elements form a group and  $\varphi_A$  is also multiplicative.

To prove the reversed statement, let  $S \subseteq \mathbb{R}$  be a subfield and  $\varphi : S \rightarrow \mathbb{R}$  be an injective field-homomorphism. Consider  $X$  as a vector space over  $S$  and let  $\{x_\gamma \mid \gamma \in \Gamma\}$  be a Hamel base of  $X$  over  $S$ . In addition, let  $\{a_\gamma \mid \gamma \in \Gamma\}$  be an arbitrary family of real numbers such that at least one of these elements is different from zero. Given an element  $x \in X$ , it can uniquely be written in the form

$$x = s_1x_{\gamma_1} + \dots + s_mx_{\gamma_m}, \tag{2.4}$$

where  $m \in \mathbb{N} \cup \{0\}$ ,  $s_1, \dots, s_m \in S$ , and  $\gamma_1, \dots, \gamma_m$  are pairwise distinct elements of the index set  $\Gamma$ . Now define  $A(x)$  by

$$A(x) := \varphi(s_1)a_{\gamma_1} + \dots + \varphi(s_m)a_{\gamma_m}.$$

Using the additivity of  $\varphi$ , it is immediate to see that  $A$  is a nonzero additive function. It remains to show that, for all  $s \in S$ ,  $(s, \varphi(s)) \in H_A$ , i.e.,

$$A(sx) = \varphi(s)A(x) \quad (x \in X). \tag{2.5}$$

If  $x$  is of the form (2.4), then  $sx = (ss_1)x_{\gamma_1} + \dots + (ss_m)x_{\gamma_m}$  and hence, by the multiplicativity of  $\varphi$ , we get

$$\begin{aligned} A(sx) &= \varphi(ss_1)a_{\gamma_1} + \dots + \varphi(ss_m)a_{\gamma_m} = \varphi(s)(\varphi(s_1)a_{\gamma_1} + \dots + \varphi(s_m)a_{\gamma_m}) = \\ &= \varphi(s)A(x), \end{aligned}$$

which completes the proof of (2.5). □

**Remark 2.4.** The equality stated in (2.3) can be rewritten as the following identity:

$$A(sx) = \varphi_A(s)A(x) \quad (s \in S_A, x \in X). \tag{2.6}$$

The additive and multiplicative properties of  $\varphi_A$  imply that if  $s \in S$  is an algebraic number over a subfield of  $\mathbb{R}$  then  $\varphi_A(s)$  must be one of its algebraic conjugates. In particular, if  $s$  is a rational number then,  $\varphi_A(s) = s$ . On the other hand, if  $s \in S$  is transcendental, then  $\varphi_A(s)$  can be any transcendental number. For an account of such results see the paper [8] by Z. Daróczy. Those real numbers  $s$  such that  $(s, s) \in H_A$  also form a subfield of  $\mathbb{R}$  (cf. Rätz [20]). This easily follows from the fact that they are characterized by the fixed point equation  $\varphi_A(s) = s$ .

An easy consequence of Theorem 2.2 and Theorem 2.3 is the following result.

**Theorem 2.5.** *Let  $T$  be a nonempty set, and  $\alpha, \beta, a, b : T \rightarrow \mathbb{R}$  be given functions satisfying property (1.3) for some  $t_0 \in T$ . Let furthermore,  $\emptyset \neq D \subset X$  be an open connected and  $(\alpha, \beta)$ -convex set. Then  $f : D \rightarrow \mathbb{R}$  is a nonconstant  $(\alpha, \beta, a, b)$ -affine function if, and only if, there exist a nonzero additive function  $A : X \rightarrow \mathbb{R}$  and a*

constant  $c \in \mathbb{R}$  such that  $\alpha(T) \cup \beta(T)$  is contained by the homogeneity field  $S_A$  of  $A$  and

$$\begin{aligned} a(t) &= \varphi_A(\alpha(t)) & (t \in T), \\ b(t) &= \varphi_A(\beta(t)) & (t \in T), \\ c(a(t) + b(t) - 1) &= 0 & (t \in T), \quad \text{and} \\ f(x) &= A(x) + c & (x \in D) \end{aligned} \quad (2.7)$$

where  $\varphi_A : S_A \rightarrow \mathbb{R}$  is the homogeneity field-homomorphism of  $A$ .

*Proof.* Applying Theorem 2.2 with  $\gamma := \alpha(t_0)$ ,  $\delta := \beta(t_0)$ ,  $p := a(t_0)$ , and  $q := b(t_0)$ , it follows that there exist an additive function  $A : X \rightarrow \mathbb{R}$  and a constant  $c \in \mathbb{R}$  such that  $f(x) = A(x) + c$  for all  $x \in D$ .

To see that the first three equations in (2.7) are valid, we substitute this form of  $f$  into (1.2) and get that, for all  $x, y \in D$  and  $t \in T$ ,

$$A(\alpha(t)x) - a(t)A(x) + A(\beta(t)y) - b(t)A(y) - c(a(t) + b(t) - 1) = 0. \quad (2.8)$$

In other words, for all fixed  $y \in D$  and  $t \in T$ , the polynomial function

$$x \mapsto A(\alpha(t)x) - a(t)A(x) + A(\beta(t)y) - b(t)A(y) - c(a(t) + b(t) - 1) \quad (x \in X)$$

vanishes on the open set  $D$ , therefore it vanishes everywhere on  $X$ . (See Székelyhidi [23].) Analogously, for all fixed  $x \in X$  and  $t \in T$ , the polynomial function

$$y \mapsto A(\alpha(t)x) - a(t)A(x) + A(\beta(t)y) - b(t)A(y) - c(a(t) + b(t) - 1) \quad (y \in X)$$

vanishes on  $D$ , therefore it vanishes everywhere on  $X$ . Therefore, (2.8) holds for all  $x, y \in X$  and  $t \in T$ .

Thus, with simple substitutions, for all  $t \in T$  and  $x \in X$ , we obtain that

$$A(\alpha(t)x) = a(t)A(x), \quad A(\beta(t)x) = b(t)A(x), \quad c(a(t) + b(t) - 1) = 0.$$

The first two equalities yield that  $(\alpha(t), a(t))$  and  $(\beta(t), b(t))$  belong to  $H_A$  for all  $t \in T$ . Therefore,  $\alpha(T) \cup \beta(T) \subseteq S_A$  and the first two equations in (2.7) are also satisfied.  $\square$

### 3. REMARKS AND EASY CONSEQUENCES OF THEOREM 2.5

**Remark 3.1.** Suppose that  $\alpha, \beta, a, b : T \rightarrow \mathbb{R}$  are given functions,  $\emptyset \neq D \subset X$  such that, for some  $t \in T$ ,

$$\alpha(t) + \beta(t) = a(t) + b(t) = 1, \quad a(t) > 0, \quad b(t) > 0, \quad \text{and} \quad \alpha(t)x + \beta(t)y, \quad \frac{x+y}{2} \in D$$

whenever  $x, y \in D$ . Then every  $(\alpha, \beta, a, b)$ -convex function  $f : D \rightarrow \mathbb{R}$  is Jensen convex, i.e.

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2} \quad (x, y \in D),$$

and every  $(\alpha, \beta, a, b)$ -affine function  $f : D \rightarrow \mathbb{R}$  satisfies the Jensen equation

$$f\left(\frac{x+y}{2}\right) = \frac{f(x) + f(y)}{2} \quad (x, y \in D).$$

In Kuczma [13, p. 315], there is an extension theorem for the Jensen equation. There  $D$  is a subset of  $\mathbb{R}^n$  with nonempty interior. Our statements follow easily from the identity (see Daróczy-Páles [9], and also Matkowski-Pycia [16])

$$\frac{x+y}{2} = \alpha(t) \left[ \alpha(t) \frac{x+y}{2} + \beta(t)y \right] + \beta(t) \left[ \alpha(t)x + \beta(t) \frac{x+y}{2} \right] \quad (x, y \in D).$$

Finally, we list some easy consequences of Theorem 2.5.

**Corollary 3.2.** *If  $\alpha(T) \cup \beta(T)$  contains a set of positive Lebesgue measure then the additive function  $A$  in Theorem 2.5 is a linear functional on  $X$  and  $a = \alpha, b = \beta$ .*

*Proof.* In this case, by a well-known theorem of Steinhaus [22], the homogeneity field  $S_A$  must contain an interval of positive length. Therefore  $S_A = \mathbb{R}$ . Thus, by the classical theorem of Darboux [7] and taking into consideration (1.3) to hold for some  $t_0 \in T$ , we have that  $\varphi_A(t) = t$  for all  $t \in \mathbb{R}$ . The remaining statements are obvious.  $\square$

The following corollary is a trivial consequence of Corollary 3.2.

**Corollary 3.3.** *Suppose that, for  $f : D \rightarrow \mathbb{R}$  and for all  $x, y \in D$ , the equality holds in the defining inequality of Breckner-convexity or Orlicz-convexity. Then  $f$  must be the constant function except the case  $s = 1$ .*

Taking into consideration Remark 2.4 (see also Daróczy [8]), we have

**Corollary 3.4.** *If  $t, q \in ]0, 1[$  are fixed,  $T = \{t\}$ ,  $\alpha(t) = t$ ,  $\beta(t) = 1 - t$ ,  $a(t) = q$ ,  $b(t) = 1 - q$  then there exists nonconstant  $(\alpha, \beta, a, b)$ -affine function if, and only if,  $t$  and  $q$  are conjugate, i.e., they are both transcendental or they are both algebraic and have the same minimal polynomial with rational coefficients.*

### Acknowledgments

*This research has been supported by the Hungarian Scientific Research Fund (OTKA) Grant NK81402 and by the TÁMOP 4.2.1./B-09/1/KONV-2010-0007 project implemented through the New Hungary Development Plan co-financed by the European Social Fund, and the European Regional Development Fund.*

## REFERENCES

- [1] P. Burai, A. Háy, *On Orlicz-convex functions*, Proc. of the 12th Symposium of Mathematics and Its Applications, Editura Politechnica, Temesvár, (2010), 73–79.
- [2] P. Burai, A. Háy, *Bernstein-Doetsch type results for generalized convex functions*, Proc. of the 12th Symposium of Mathematics and Its Applications, Editura Politechnica, Temesvár, (2010), 118–124.
- [3] P. Burai, A. Háy, T. Juhász, *Bernstein-Doetsch type results for  $s$ -convex functions*, Publ. Math. Debrecen **75** (2009) 1–2, 23–31.
- [4] P. Burai, A. Háy, *On approximately  $h$ -convex functions*, accepted for publication, Journal of Convex Analysis, available electronically <http://www.heldermann.de/JCA/JCA18/JCA182/jca18029.htm>.
- [5] W.W. Breckner, *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen*, Publ. Inst. Math. (Beograd) **23** (1978), 13–20.
- [6] W.W. Breckner, *Hölder-continuity of certain generalized convex functions*, Optimization **28** (1994), 201–209.
- [7] G. Darboux, *Sur la composition des forces en statique*, Bull. Sci. Math. **9** (1875) 1, 281–288.
- [8] Z. Daróczy, *Notwendige und hinreichende Bedingungen für die Existenz von nichtkonstanten Lösungen linearer Funktionalgleichungen*, Acta Sci. Math. (Szeged) **22** (1961), 31–41.
- [9] Z. Daróczy, Zs. Páles, *Convexity with given infinite weight sequences*, Stochastica **11** (1987), 5–12.
- [10] E.K. Godunova, V.I. Levin, *Neravenstva dlja funkcii širokogo klassa, soderzascego vy-puklye, monotonnnye i nekotorye drugie vidy funkii*, Vycislitel. Mat. i. Fiz. Mezvuzov. Sb. Nauc. Trudov, MGPI, Moskva, 1985, pp. 138–142.
- [11] A. Háy, *Bernstein-Doetsch type results for  $h$ -convex functions*, accepted for publication, Math. Ineq. Appl. (2011).
- [12] H. Hudzik, L. Maligranda, *Some remarks on  $s_i$ -convex functions*, Aequationes Math. **48** (1994), 100–111.
- [13] M. Kuczma, *An Introduction to the Theory of Functional Equations and Inequalities*, Prace Naukowe Uniwersytetu Śląskiego w Katowicach Vol. CDLXXXIX (Państwowe Wydawnictwo Naukowe – Uniwersytet Śląski, Warszawa–Kraków–Katowice, 1985).
- [14] N. Kuhn, *A note on  $t$ -convex functions*, General Inequalities 4, Internat. Ser. Numer. Math. **71** (1984), 269–276.
- [15] N. Kuhn, *On the structure of  $(s, t)$ -convex functions*, General Inequalities 5, Internat. Ser. Numer. Math. **80** (1987), 161–174.
- [16] J. Matkowski, M. Pycia, *On  $(\alpha, a)$ -convex functions*, Arch. Math (Basel) **64** (1995), 132–138.
- [17] W. Orlicz, *A note on modular spaces I*, Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys. **9** (1961), 157–162.



- [18] C.E.M. Pearce, A.M. Rubinov, *P-functions, quasi-convex functions and Hadamard-type inequalities*, J. Math. Anal. Appl. **240** (1999), 92–104.
- [19] F. Radó, J.A. Baker, *Peirder's equation and aggregation of allocations*, Aequationes Math. **32** (1987), 227–239.
- [20] J. Rätz, *On the homogeneity of additive mappings*, Aequationes Math. **14** (1976), 67–71.
- [21] A.W. Roberts, D.E. Varberg, *Convex Functions*, Academic Press, New York, 1973.
- [22] H. Steinhaus, *Sur les distances des points des ensembles de mesure positive*, Fund. Math. **1** (1920), 99–104.
- [23] L. Székelyhidi, *Regularity properties of polynomials on groups*, Acta Math. Hung. **45** (1985), 15–19.
- [24] S. Varošanec, *On  $h$ -convexity*, J. Math. Anal. Appl. **32** (2007), 303–311.

Gyula Maksa  
maksa@math.klte.hu

University of Debrecen  
Institute of Mathematics  
H-4010 Debrecen, Pf. 12, Hungary

Zsolt Páles  
pales@math.klte.hu

University of Debrecen  
Institute of Mathematics  
H-4010 Debrecen, Pf. 12, Hungary

*Received: October 20, 2010.*

*Accepted: October 28, 2010.*