

## ON THE CHROMATIC NUMBER OF $(P_5, \text{windmill})$ -FREE GRAPHS

Ingo Schiermeyer

*Communicated by Hao Li*

**Abstract.** In this paper we study the chromatic number of  $(P_5, \text{windmill})$ -free graphs. For integers  $r, p \geq 2$  the *windmill graph*  $W_{r+1}^p = K_1 \vee pK_r$  is the graph obtained by joining a single vertex (the center) to the vertices of  $p$  disjoint copies of a complete graph  $K_r$ . Our main result is that every  $(P_5, \text{windmill})$ -free graph  $G$  admits a polynomial  $\chi$ -binding function. Moreover, we will present polynomial  $\chi$ -binding functions for several other subclasses of  $P_5$ -free graphs.

**Keywords:** vertex colouring, perfect graphs,  $\chi$ -binding function, forbidden induced subgraph.

**Mathematics Subject Classification:** 05C15, 05C17.

### 1. INTRODUCTION

We consider finite, simple, and undirected graphs, and use standard terminology and notation.

Let  $G$  be a graph. An *induced subgraph* of  $G$  is a graph  $H$  such that  $V(H) \subseteq V(G)$ , and  $uv \in E(H)$  if and only if  $uv \in E(G)$  for all  $u, v \in V(H)$ . Given graphs  $G$  and  $F$  we say that  $G$  *contains*  $F$  if  $F$  is isomorphic to an induced subgraph of  $G$ . We say that a graph  $G$  is  $F$ -free, if it does not contain  $F$ . For two graphs  $G, H$  we denote by  $G + H$  the disjoint union and by  $G \vee H$  the join of  $G$  and  $H$ , respectively.

A graph  $G$  is called  $k$ -colourable, if its vertices can be coloured with  $k$  colours so that adjacent vertices obtain distinct colours. The smallest  $k$  such that a given graph  $G$  is  $k$ -colourable is called its *chromatic number*, denoted by  $\chi(G)$ . It is well-known that  $\omega(G) \leq \chi(G) \leq \Delta(G) + 1$  for any graph  $G$ , where  $\omega(G)$  denotes its clique number and  $\Delta(G)$  its maximum degree. A graph  $G$  is *perfect* if  $\chi(H) = \omega(H)$  for every induced subgraph  $H$  of  $G$ .

A family  $\mathcal{G}$  of graphs is called  $\chi$ -bound with binding function  $f$  if  $\chi(G') \leq f(\omega(G'))$  holds whenever  $G \in \mathcal{G}$  and  $G'$  is an induced subgraph of  $G$ . For a fixed graph  $H$  let  $\mathcal{G}(H)$  denote the family of graphs which are  $H$ -free.

The following theorems are well known in chromatic graph theory.

**Theorem 1.1** (Erdős [5]). *For any positive integers  $k, l \geq 3$  there exists a graph  $G$  with girth  $g(G) \geq l$  and chromatic number  $\chi(G) \geq k$ .*

**Theorem 1.2** (The Strong Perfect Graph Theorem [4]). *A graph is perfect if and only if it contains neither an induced odd cycle of length at least five nor its complement.*

In this paper we study the chromatic number of  $P_5$ -free graphs. Our work was motivated by the following conjecture of Gyárfás.

**Conjecture 1.3** (Gyárfás' Conjecture [8]). *Let  $T$  be any tree (or forest). Then there is a function  $f_T$  such that every  $T$ -free graph  $G$  satisfies  $\chi(G) \leq f_T(\omega(G))$ .*

Gyárfás [8] proved this conjecture when  $T$  is a path  $P_k$  for all  $k \geq 4$  by showing

**Theorem 1.4.** *Let  $G$  be a  $P_k$ -free graph for  $k \geq 4$  with clique number  $\omega(G) \geq 2$ . Then*

$$\frac{R(\omega + 1, \lceil \frac{k}{2} \rceil) - 1}{\lceil \frac{k}{2} \rceil - 1} \leq f_k(\omega) \leq (k - 1)^{\omega(G) - 1},$$

where  $R(s, t)$  is the Ramsey number.

Note that  $P_4$ -free graphs are perfect. The currently best known upper bound for  $P_5$ -free graphs is due to Esperet, Lemoine, Maffray, and Morel [6].

**Theorem 1.5.** *Let  $G$  be a  $P_5$ -free graph with clique number  $\omega(G) \geq 3$ . Then  $\chi(G) \leq 5 \cdot 3^{\omega(G) - 3}$ .*

One may wonder whether this exponential bound can be improved. In particular:

**Question 1.6.** *Are there polynomial ( $\chi$ -binding) functions  $f_k$  for  $k \geq 5$  such that every  $P_k$ -free graph  $G$  satisfies  $\chi(G) \leq f_k(\omega(G))$ ?*

If there would be a polynomial ( $\chi$ -binding) function  $f_k$  for some  $k \geq 5$ , then it would imply the Erdős-Hajnal conjecture for  $P_k$ -free graphs. The Erdős-Hajnal conjecture states that for every graph  $H$ , there exists a constant  $\delta(H) > 0$  such that every graph  $G$  with no induced subgraph isomorphic to  $H$  has either a clique or a stable set of size at least  $|V(G)|^{\delta(H)}$ . However, the Erdős-Hajnal conjecture is still open for  $P_k$ -free graphs for all  $k \geq 5$  (cf. [3] for a survey).

## 2. GENERAL GRAPHS

One of the earliest results is due to Wagon, who has considered graphs without induced matchings.

**Theorem 2.1** ([10]). *Let  $G$  be a  $2K_2$ -free graph with clique number  $\omega(G)$ . Then  $\chi(G) \leq \binom{\omega(G)+1}{2}$ .*

This theorem admits a nice generalization as follows.

**Theorem 2.2.** *Let  $H$  be a graph such that  $\mathcal{G}(H)$  has an  $O(\omega^t)$   $\chi$ -binding function for some  $t \geq 1$ , and let  $G$  be a  $K_2 + H$ -free graph with clique number  $\omega(G)$ . Then  $G$  has an  $O(\omega^{2+t})$   $\chi$ -binding function.*

The proof for Theorem 2.2 will be given after the proof of Theorem 3.6.

**Theorem 2.3** ([10]). *The family  $\mathcal{G}(pK_2)$  has an  $O(\omega^{2p-2})$   $\chi$ -binding function for all  $p \geq 1$ .*


Note that the statement of Theorem 2.3 can be made more precise as follows. In [10] a  $\chi$ -binding function  $f_p(\omega)$  for the class of  $pK_2$ -free graphs was defined by  $f_1(\omega) = 1, f_{p+1}(\omega) = \binom{\omega}{2} f_p(\omega) + \omega$ . From this one can deduce that  $f_p(\omega) \leq (\omega+1) \frac{\omega^{2p-3}}{2^{p-1}}$  for all  $p \geq 1$ .

Like Theorem 2.2 for Theorem 2.1, Theorem 2.3 has the following counterpart.


**Theorem 2.4.** *Let  $H$  be a graph such that  $\mathcal{G}(H)$  has an  $O(\omega^t)$   $\chi$ -binding function for some  $t \geq 1$ , and let  $G$  be a  $pK_2 + H$ -free graph with clique number  $\omega(G)$ . Then  $G$  has an  $O(\omega^{2p-2+t})$   $\chi$ -binding function.*

### 3. $P_k$ -FREE GRAPHS

In this section we will consider  $P_k$ -free graphs. Since  $P_4$ -free graphs are perfect graphs, we may assume  $k \geq 5$ . For the presentation of our results we will need several forbidden induced subgraphs, which are presented in Figure 1 and Figure 2. The following results have been shown for  $P_5$ -free graphs.



**Fig. 1.** The graphs *House*, *Paw*, *Diamond*, and *Gem*.



**Fig. 2.** The graphs *Claw*, *Dart*, *Cricket*,  $Gem^+$ , and *Windmill*  $W_3^2$

**Theorem 3.1** ([7]). *Let  $G$  be a connected  $(P_5, \text{House})$ -free graph of order  $n$  and clique number  $\omega(G)$ . Then  $\chi(G) \leq \binom{\omega(G)+1}{2}$ .*

**Theorem 3.2** ([2]). *Let  $G$  be a connected  $(P_5, \text{Gem})$ -free graph of order  $n$  and clique number  $\omega(G)$ . Then  $\chi(G) \leq 6\omega(G)$ .*

Since the Paw and the Diamond are both induced subgraphs of the Gem, we obtain the following corollary.

**Corollary 3.3.** *Let  $G$  be a connected  $(P_5, H)$ -free graph of order  $n$  and clique number  $\omega(G)$ , where  $H \in \{\text{Paw}, \text{Diamond}\}$ . Then  $\chi(G) \leq 6\omega(G)$ .*

In [9] the subgraph Gem was replaced by the supergraph  $\text{Gem}^+ = K_1 \vee (K_1 + P_4)$ .

**Theorem 3.4.** *Let  $G$  be a  $(P_5, \text{Gem}^+)$ -free graph of order  $n$  and clique number  $\omega(G)$ . Then  $\chi(G) \leq \omega^2(G)$ .*

Since the Claw, the Dart and the Cricket are induced subgraphs of the  $\text{Gem}^+$ , we obtain the following corollary.

**Corollary 3.5.** *Let  $G$  be a connected  $(P_5, H)$ -free graph of order  $n$  and clique number  $\omega(G)$ , where  $H \in \{\text{Claw}, \text{Dart}, \text{Cricket}\}$ . Then  $\chi(G) \leq \omega^2(G)$ .*

We start by proving a generalization of Theorem 2.1 for  $P_5$ -free graphs.

**Theorem 3.6.** *Let  $G$  be a  $(P_k, K_{n_1} + K_{n_2})$ -free graph for some  $n_1 \geq n_2 \geq 2$ . Then  $\chi(G) \leq c(n_1) \cdot \omega^{n_1}$  for a constant  $c(n_1)$ .*

*Proof.* Let  $\omega = \omega(G)$  and let  $F$  be a complete subgraph of  $G$  with  $|V(F)| = \omega$ . For a subset  $T \subset V(F)$  with  $|T| = t$  and  $1 \leq t \leq n_1 - 1$ , let

$$M(T) = \{v \in V(G) \setminus F \mid N(v) \cap V(F) = V(F) \setminus T\}.$$

Then  $G[M(T)]$  is  $K_{t+1}$ -free, since otherwise there would be a complete subgraph of  $G$  of order at least  $(\omega - t) + (t + 1) = \omega + 1$ , a contradiction. Hence  $\chi(G[M(T)]) \leq f_{P_k}(t)$ . For a subset  $T \subset V(F)$  with  $|T| = n_1$  let

$$M(T) = \{v \in V(G) \setminus F \mid N(v) \cap V(F) \subseteq V(F) \setminus T\}.$$

Then  $G[M(T)]$  is  $K_{n_2}$ -free, since  $G$  is  $(K_{n_1} + K_{n_2})$ -free. Hence we have  $\chi(G[M(T)]) \leq f_{P_5}(n_2 - 1)$ .

We now colour the vertices of  $G$  as follows. The vertices of  $F$  obtain  $\omega$  distinct colours. For every vertex  $w \in V(F)$ , the set  $w \cup M(\{w\})$  is independent. So all vertices of  $M(\{w\})$  obtain the same colour as  $w$ . Next for every subset  $T \subset V(F)$  with  $2 \leq t \leq n_1$  we choose a private set of  $f_{P_k}(t)$  colours. For every  $t$  with  $2 \leq t \leq n_1$  there are  $\binom{\omega}{t}$  subsets  $T \subset V(F)$  with  $|T| = t$ . So we obtain

$$\chi(G) \leq \omega + \sum_{t=2}^{n_1} \binom{\omega}{t} f_{P_k}(t),$$

which is a polynomial of degree  $n_1$  in  $\omega$ . □

*Proof of Theorem 2.2.* We can follow the proof of Theorem 3.6 with the following modification. With  $n_1 = 2$  and  $K_{n_2}$  replaced by  $H$  we obtain

$$\chi(G) \leq \omega + \binom{\omega}{2} O(\omega^t).$$

Hence  $G$  has an  $O(\omega^{2+t})$   $\chi$ -binding function. □

Theorem 3.6 can be generalized to  $(P_k, K_{n_1} + K_{n_2} + \dots + K_{n_p})$ -free graphs for  $p \geq 3$  and  $n_1 \geq n_2 \geq \dots \geq n_p$  as follows. We use the proof above as an induction step with the following modification. For a subset  $T \subset V(F)$  with  $|T| = n_1$  the subgraph  $G[M(T)]$  is  $(P_k, K_{n_2} + \dots + K_{n_p})$ -free. Hence

$$\chi(G[M(T)]) \leq c(n_2, \dots, n_p) \cdot \omega^{\sum_{i=2}^{p-1} n_i},$$

which leads to

$$\chi(G) \leq \omega + \sum_{t=2}^{n_1-1} \binom{\omega}{t} f_{P_k}(t) + \binom{\omega}{n_1} \cdot c(n_2, \dots, n_p) \cdot \omega^{\sum_{i=2}^{p-1} n_i},$$

which is a polynomial of degree  $\sum_{i=1}^{p-1} n_i$  in  $\omega$ . So we obtain the following result.

**Theorem 3.7.** *Let  $G$  be a  $(P_k, K_{n_1} + K_{n_2} + \dots + K_{n_p})$ -free graph for some  $p \geq 2$  and  $n_1 \geq n_2 \geq \dots \geq n_p \geq 2$ . Then*

$$\chi(G) \leq c(n_1, \dots, n_p) \cdot \omega^{\sum_{i=1}^{p-1} n_i}$$

for a constant  $c(n_1, \dots, n_p)$ .

Theorem 3.6 also leads to the following variation.

**Theorem 3.8.** *Let  $G$  be a  $(P_k, K_{n_1} + P_4)$ -free graph for some  $n_1 \geq 2$ . Then  $\chi(G) \leq c(n_1) \cdot \omega^{n_1+1}$  for a constant  $c(n_1)$ .*

*Proof.* We follow the proof of Theorem 3.6 with the following modification: For a subset  $T \subset V(F)$  with  $|T| = n_1$  let

$$M(T) = \{v \in V(G) \setminus F \mid N(v) \cap V(F) \subseteq V(F) \setminus T\}.$$

Then  $G[M(T)]$  is  $P_4$ -free, since  $G$  is  $(K_{n_1} + P_4)$ -free. Hence  $\chi(G[M(T)]) \leq \omega$ , since  $P_4$ -free graphs are perfect graphs. So we obtain

$$\chi(G) \leq \omega + \sum_{t=2}^{n_1-1} \binom{\omega}{t} f_{P_k}(t) + \binom{\omega}{n_1} \omega,$$

which is a polynomial of degree  $n_1 + 1$  in  $\omega$ . □

The counterpart of Theorem 2.2 for  $P_k$ -free graphs is the following.

**Theorem 3.9.** *Let  $H$  be a graph such that  $\mathcal{G}(H)$  has an  $O(\omega^t)$   $\chi$ -binding function for some  $t \geq 1$ , and let  $G$  be a  $(P_k, K_{n_1} + H)$ -free graph for some  $n_1 \geq 2$ . Then  $\chi(G) \leq c(n_1, H) \cdot \omega^{n_1+t}$  for a constant  $c(n_1, H)$ .*

4.  $(P_5, \text{windmill})$ -FREE GRAPHS

For integers  $r, p \geq 2$  the *windmill graph*  $W_{r+1}^p = K_1 \vee pK_r$  is the graph obtained by joining a single vertex (the center) to the vertices of  $p$  disjoint copies of a complete graph  $K_r$  (the Windmill  $W_3^2$  is shown in Figure 2). For integers  $n_1 \geq n_2 \geq \dots \geq n_p \geq 2$ , the *generalized windmill graph*  $W(n_1, n_2, \dots, n_p) = K_1 \vee (K_{n_1} + K_{n_2} + \dots + K_{n_p})$  is the graph obtained by joining a single vertex (the center) to the vertices of  $p$  disjoint complete graphs  $K_{n_1}, \dots, K_{n_p}$ .

We start with a structural result for connected  $P_5$ -free graphs.

**Theorem 4.1** (Bacsó and Tuza [1]). *Every connected  $P_5$ -free graph contains a dominating clique or a dominating  $P_3$ .*

This admits the following result for  $P_5$ -free graphs.

**Theorem 4.2.** *Let  $H$  be a graph such that  $\mathcal{G}(H)$  has an  $O(\omega^t)$   $\chi$ -binding function for some  $t \geq 1$ , and let  $G$  be a connected  $(P_5, K_1 \vee H)$ -free graph with clique number  $\omega(G)$ . Then  $G$  has an  $O(\omega^{t+1})$   $\chi$ -binding function.*

*Proof.* Let  $D = \{w_1, w_2, \dots, w_d\}$  be a dominating set as in Theorem 4.1 with  $d = |D|$ . We may assume  $\omega \geq 3$ , since otherwise  $\chi(G) \leq 3$  by Theorem 1.4. For  $1 \leq i \leq d$ , let  $G_i = \{w_i\} \cup (N(w_i) \cap (V(G) - D))$ . Then  $\chi(G) \leq \sum_{i=1}^d \chi(G_i)$ . Moreover, if  $G_i - w_i$  is  $H$ -free, then  $G_i$  is  $(K_1 \vee H)$ -free. Then  $G$  has an  $O(\omega^{t+1})$   $\chi$ -binding function, since  $H$  has an  $O(\omega^t)$   $\chi$ -binding function for some  $t \geq 1$  and  $d \leq \omega$ .  $\square$

So we can apply Theorem 4.2 to obtain the following results for  $(P_5, \text{windmill})$ -free graphs.

**Theorem 4.3.** *Let  $G$  be a  $(P_5, W(n_1, n_2))$ -free graph for some  $n_1 \geq n_2 \geq 2$ . Then  $\chi(G) \leq c(n_1) \cdot \omega^{n_1+1}$  for a constant  $c(n_1)$ .*

**Theorem 4.4.** *Let  $G$  be a  $(P_5, W(n_1, n_2, \dots, n_p))$ -free graph for some  $p \geq 2$  and  $n_1 \geq n_2 \geq \dots \geq n_p \geq 2$ . Then*

$$\chi(G) \leq c(n_1, \dots, n_p) \cdot \omega^{1 + \sum_{i=1}^{p-1} n_i}$$

for a constant  $c(n_1, \dots, n_p)$ .

## REFERENCES

- [1] G. Bacsó, Zs. Tuza, *Dominating cliques in  $P_5$ -free graphs*, Period. Math. Hungar. **21** (1990) 3, 303–308.
- [2] S.A. Choudum, T. Karthick, M.A. Shalu, *Perfect coloring and linearly  $\chi$ -bound  $P_6$ -free graphs*, J. Graph Theory **54** (2006) 4, 293–306.
- [3] M. Chudnovsky, *The Erdős-Hajnal conjecture – a survey*, J. Graph Theory **75** (2014), 178–190.

- [4] M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas, *The strong perfect graph theorem*, Ann. of Math. **164** (2006), 51–229.
- [5] P. Erdős, *Graph theory and probability*, Canad. J. Math. **11** (1959), 34–38.
- [6] L. Esperet, L. Lemoine, F. Maffray, G. Morel, *The chromatic number of  $\{P_5, K_4\}$ -free graphs*, Discrete Math. **313** (2013), 743–754.
- [7] J.L. Fouquet, V. Giakoumakis, F. Maire, H. Thuillier, *On graphs without  $P_5$  and  $\overline{P}_5$* , Discrete Math. **146** (1995), 33–44.
- [8] A. Gyárfás, *Problems from the world surrounding perfect graphs*, [in:] Proc. Int. Conf. on Comb. Analysis and Applications, Pokrzywna, 1985; Zastos. Mat. **19** (1987), 413–441.
- [9] I. Schiermeyer, *Chromatic number of  $P_5$ -free graphs: Reed's conjecture*, Discrete Math. **339** (2016) 7, 1940–1943.
- [10] S. Wagon, *A bound on the chromatic number of graphs without certain induced subgraphs*, J. Combin. Theory Ser. B **29** (1980), 345–346.

Ingo Schiermeyer  
Ingo.Schiermeyer@tu-freiberg.de

Technische Universität Bergakademie Freiberg  
Institut für Diskrete Mathematik und Algebra  
09596 Freiberg, Germany

*Received: July 25, 2016.*

*Accepted: November 9, 2016.*