

DAMPED NONLINEAR GINZBURG–LANDAU EQUATION WITH SATURATION. PART II. STRONG STABILIZATION

Pascal Bégout and Jesús Ildefonso Díaz

Communicated by Vicențiu D. Rădulescu

Abstract. We study the complex Ginzburg–Landau equation posed on possibly unbounded domains, including some singular and saturated nonlinear damping terms. This model interpolates between the nonlinear Schrödinger equation and dissipative parabolic dynamics through a complex time-derivative prefactor, capturing the interplay between dispersion and dissipation. As a continuation of our previous study on the existence and uniqueness of solutions, we prove here some strong stabilization properties. In particular, we show the finite time extinction of solutions induced by the nonlinear saturation mechanism, which, sometimes, can be understood as a bang-bang control. The analysis relies on refined energy methods. Our results provide a rigorous justification of nonlinear dissipation as an effective stabilization mechanism for this class of complex equations where the maximum principle fails.

Keywords: damped Ginzburg–Landau equation, saturated nonlinearity, finite time extinction.

Mathematics Subject Classification: 35Q56, 35B40, 93D40.

1. INTRODUCTION

The complex Ginzburg–Landau equation constitutes one of the most fundamental models in the theory of nonlinear dissipative systems. For a more detailed introduction to the model we will consider in this paper we send the reader to the Part I of our study (see [9]).

In several recent works (see, e.g., [8]), the strong stabilization of a damped nonlinear Schrödinger equation with saturation effects was established on unbounded domains. That analysis demonstrates that suitably chosen nonlinear damping mechanisms can overcome dispersive effects even in the absence of compactness properties typically available in bounded domains. Such results are particularly relevant for physical systems modeled in open space, where boundary confinement cannot be assumed. The main goal of this paper is to extend the general approach taken in the theory presented in [8] in order to extend previous results in the literature on complex Ginzburg–Landau

equation in which the saturation term is understood as an absorption term (see, e.g., Antontsev, Dias and Figueira [1] and [11–15]).

The damped nonlinear Schrödinger equation may be viewed as a limiting or simplified model within the broader Ginzburg–Landau framework. Introducing a complex coefficient in front of the time derivative allows one to interpolate continuously between purely dispersive Schrödinger dynamics and purely dissipative parabolic dynamics. This observation motivates the extension of the stabilization theory developed in [8] to the complex Ginzburg–Landau equation posed on general domains $\Omega \subseteq \mathbb{R}^N$ (possibly unbounded), with boundary $\partial\Omega$,

$$\begin{cases} e^{-i\theta} \frac{\partial u}{\partial t} - \Delta u + a|u|^{-(1-m)}u + b|u|^{p-1}u + \gamma u = f & \text{in } (0, \infty) \times \Omega, & (1.1) \\ u|_{\partial\Omega} = 0 & \text{on } (0, \infty) \times \partial\Omega, & (1.2) \\ u(0) = u_0 & \text{in } \Omega, & (1.3) \end{cases}$$

where $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $0 \leq m \leq 1$ and $a, b, \gamma \in \mathbb{C}$. Here we write, for generality $p \in (1, \infty)$ but the physically more often case considered in the literature corresponds to $p = 3$.

For $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and $m \geq 0$, we introduce the following set of complex numbers:

$$C_\theta(m) = \left\{ z \in \mathbb{C}; \operatorname{Re}(ze^{i\theta}) > 0 \text{ and } 2\sqrt{m} \operatorname{Re}(ze^{i\theta}) \geq |1 - m| |\operatorname{Im}(ze^{i\theta})| \right\}.$$

In the particular cases in which $m \in \{0, 1\}$, the set $C_\theta(m)$ becomes

$$\begin{aligned} C_\theta(0) &= \left\{ z \in \mathbb{C}; \operatorname{Re}(ze^{i\theta}) > 0 \text{ and } \operatorname{Im}(ze^{i\theta}) = 0 \right\}, \\ C_\theta(1) &= \left\{ z \in \mathbb{C}; \operatorname{Re}(ze^{i\theta}) > 0 \right\}, \end{aligned}$$

and actually

$$C_\theta(0) = \left\{ z \in \mathbb{C}; \text{ there exists } \mu > 0 \text{ such that } z = \mu e^{-i\theta} \right\}.$$

We note that if $\theta = \frac{\pi}{2}$, $0 \leq m \leq 1$, $a \in C_\theta(m)$, $b = 0$, $\gamma = -V(x) \in L^1_{\text{loc}}(\Omega; \mathbb{R})$ and $f \in L^1_{\text{loc}}([0, \infty); L^2(\Omega))$, then equation (1.1) becomes

$$i \frac{\partial u}{\partial t} + \Delta u + V(x)u - a|u|^{-(1-m)}u = -f. \quad (1.4)$$

It follows that the nonlinear Schrödinger equation (1.4) is a limit case of the Ginzburg–Landau equation (1.1), in term of θ . But the Ginzburg–Landau equation (1.1) may also be considered as an intermediate equation between the nonlinear Schrödinger equation and the nonlinear heat equation

$$\frac{\partial u}{\partial t} - \Delta u + a|u|^{-(1-m)}u = f,$$

by taking $\theta = 0$, $a \in \mathbb{R}$ and $b = \gamma = 0$ in (1.1). In this last case, $a \in C_0(m)$ only means that a is a positive real number.

The strategy of the proofs in this paper relies on the use of suitable energy methods, sharpening the ones presented in the monograph [2]. Those methods capture the effective dissipation induced by the nonlinear terms, combined with refined energy estimates adapted to unbounded domains. Singular nonlinearities with $0 \leq m < 1$ require weak formulations and truncation arguments to control the dynamics near vanishing amplitudes.

In stabilization problems, the presence of a damped saturated term plays a crucial role. Linear mechanisms and Lipschitz nonlinear terms alone produce only exponential decay but not strong stabilization properties.

From a physical standpoint, the stabilization mechanism analyzed in this work can be interpreted as an effective dissipation process capable of counterbalancing dispersion and diffusion in open systems. In unbounded spatial domains, energy injected locally can escape to infinity through wave propagation or diffusive transport, preventing the formation of confined modes and undermining stabilization mechanisms based solely on linear damping.

The nonlinear terms appearing in the complex Ginzburg–Landau equation introduce amplitude-dependent dissipation that becomes particularly effective in regimes where linear mechanisms fail. The singular term $|u|^{-(1-m)}u$ acts as a strong damping mechanism near low-amplitude states, suppressing residual oscillations and preventing the persistence of small-amplitude coherent structures. From the physical point of view, this term can be interpreted as a saturation or threshold effect that inhibits the survival of weak excitations.

The complex prefactor $e^{-i\theta}$ plays a fundamental role in shaping the dynamics. For $\theta \neq 0$, the system no longer conserves energy in the Hamiltonian sense, and the interaction between dispersive and dissipative components leads to a gradual relaxation toward equilibrium. This behavior is characteristic of systems far from equilibrium, where dissipation and dispersion coexist and compete.

From the perspective of nonlinear dynamics, the stabilization results obtained in this work indicate that the complex Ginzburg–Landau equation on unbounded domains behaves as a genuinely dissipative system, despite the absence of geometric confinement and the presence of continuous spectrum. The nonlinear damping mechanisms effectively restore asymptotic stability by suppressing long-wavelength excitations and dispersive tails, leading to strong convergence toward stationary states.

These results provide a rigorous mathematical justification for the physical intuition that nonlinear dissipation and saturation can stabilize extended systems even in open geometries, a phenomenon observed in a variety of physical contexts ranging from superconductivity and nonlinear optics to pattern-forming systems far from equilibrium.

One of our main motivations is the rigorous proof of the strong stabilization (in a finite time) to $u = 0$. This qualitative property is also called in the literature as the Finite Time Extinction property, and it is also related with the so-called Finite Time Null controllability in Control Theory. For instance, the case of a pure saturation $m = 0$ nonlinearity, as the one considered in (1.1) can be understood also in the framework of Control Theory as a special case of a feed-back control $y(t, x)$ of “bang-bang type” for the complex Ginzburg–Landau equation when we write (1.1) in the form

$$e^{-i\theta} \frac{\partial u}{\partial t} - \Delta u + b|u|^{p-1}u + \gamma u = f + y(t, x), \quad \text{in } (0, \infty) \times \Omega,$$

with

$$y(t, x) = -i\mu \frac{u(t, x)}{|u(t, x)|}.$$

where $\mu > 0$. This type of control has been considered in the applications to many dissipative evolution equations (see [8] and its references). Nevertheless, the controllability for the complex Ginzburg–Landau equation is more delicate (for some related results, see, e.g., Rosier and Zhang [20] and Fenza, Labbadi and Ouzahra [17]).

In this paper, finite time extinction property (finite stabilization) of the solutions are obtained under the assumption that $a \in C_\theta(m)$, while for the equation (1.4), they are proved in the series of papers [3, 5–8] under the assumption that $-a \in C(m)$, where

$$C(m) = \left\{ z \in \mathbb{C}; \operatorname{Im}(z) > 0 \text{ and } 2\sqrt{m}\operatorname{Im}(z) \geq (1 - m)|\operatorname{Re}(z)| \right\}.$$

Finally, notice that $a \in C_{\frac{\pi}{2}}(m)$ if, and only if, $-a \in C(m)$.

The organization of this paper is the following. Section 2 presents the statements of the main results concerning the strong stabilization. In Section 3 we present the proofs of the results concerning the strong stabilization of the solutions.

We collect here some notations that will be used along with this paper. For $t \in \mathbb{R}$, $t_+ = \max\{t, 0\}$ is the positive part of t . Unless if specified, all functions are complex-valued and all the vector spaces are considered over the field \mathbb{R} . For a Banach space X , we denote by $X^* \stackrel{\text{def}}{=} \mathcal{L}(X; \mathbb{R})$ its topological dual and by $\langle \cdot, \cdot \rangle_{X^*, X} \in \mathbb{R}$ the $X^* - X$ duality product. The product $iT \in X^*$, for $T \in X^*$, is defined in [4]. For $1 \leq p \leq \infty$, p' is the conjugate of p defined by $\frac{1}{p} + \frac{1}{p'} = 1$. For a Banach space X and $p \in (0, \infty]$, $u \in L_{\text{loc}}^p([0, \infty); X)$ means that $u \in L_{\text{loc}}^p((0, \infty); X)$ and for any $T > 0$, $u|_{(0, T)} \in L^p((0, T); X)$. In the same way, we will use the notation $u \in W_{\text{loc}}^{1, p}([0, \infty); X)$. If $p \in (0, \infty]$ and $r = 0$, then $L^{\frac{p}{r}}(\Omega) = L^\infty(\Omega)$ and $W^{1, \frac{p}{r}}(\Omega) = W^{1, \infty}(\Omega)$. Finally, we denote by C auxiliary positive constants, and sometimes, for positive parameters a_1, \dots, a_n , write as $C(a_1, \dots, a_n)$ to indicate that the constant C depends only and continuously on a_1, \dots, a_n (we will use this convention for constants which are not denoted merely by “ C ”).

2. FINITE TIME EXTINCTION PROPERTY

Let us recall that if $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and $m \geq 0$, then $C_\theta(m)$ is defined by

$$C_\theta(m) = \left\{ z \in \mathbb{C}; \operatorname{Re}(ze^{i\theta}) > 0 \text{ and } 2\sqrt{m}\operatorname{Re}(ze^{i\theta}) \geq |1 - m| |\operatorname{Im}(ze^{i\theta})| \right\}.$$

In order to have existence of solution, we make the following assumptions.

Assumption 2.1. We assume the following.

$$\begin{aligned} \Omega &\text{ is any nonempty open subset of } \mathbb{R}^N, \\ -\frac{\pi}{2} &< \theta < \frac{\pi}{2}, \\ m &\in [0, 1] \text{ and } p \in (1, \infty), \\ a &\in C_\theta(m) \text{ and } b \in C_\theta(p) \cup \{0\}, \\ \gamma &\in \mathbb{C} \text{ with } \operatorname{Re}(\gamma e^{i\theta}) \geq 0. \end{aligned}$$

Definition 2.2. Let $\mathcal{O} \subseteq \mathbb{R}^N$ be an open subset and let $u \in L^1_{\text{loc}}(\mathcal{O})$. A function U is said to be a *saturated section* associated to u if $U \in L^\infty(\mathcal{O})$, $\|U\|_{L^\infty(\mathcal{O})} \leq 1$ and $U = \frac{u}{|u|}$, almost everywhere, where $u \neq 0$.

Now, let us recall the notion of solution.

Definition 2.3. Let Assumption 2.1 be fulfilled, let $f \in L^1_{\text{loc}}([0, \infty); L^2(\Omega))$ and let $u_0 \in L^2(\Omega)$. We shall say that u is an *H^2 -solution* to (1.1)–(1.3), if u satisfies the following properties.

1. We have that

$$u \in L^{m+1}_{\text{loc}}([0, \infty); H^1_0(\Omega) \cap X_{m,p}) \cap W^{1, \frac{m+1}{m}}_{\text{loc}}([0, \infty); L^2(\Omega) + X^*_{m,p}), \quad (2.1)$$

where $X_{m,p} = L^{m+1}(\Omega) \cap L^{p+1}(\Omega)$.

2. For almost every $t > 0$, $\Delta u(t) \in L^2(\Omega)$.
3. a) If $m > 0$, then u satisfies (1.1) in $\mathcal{D}'((0, \infty) \times \Omega)$.
 b) If $m = 0$, then there exists a saturated section U associated to u such that the pair (u, U) satisfies

$$e^{-i\theta} \frac{\partial u}{\partial t} - \Delta u + aU + b|u|^{p-1}u + \gamma u = f, \quad (2.2)$$

in $\mathcal{D}'((0, \infty) \times \Omega)$.

4. We have that $u(0) = u_0$, in $L^2(\Omega)$.

We shall say that u is an *L^2 -solution* or a *weak solution* to (1.1)–(1.3) if there exists a pair

$$(u_n, f_n)_{n \in \mathbb{N}} \subset C([0, \infty); L^2(\Omega)) \times L^1_{\text{loc}}([0, \infty); L^2(\Omega)), \quad (2.3)$$

such that for any $n \in \mathbb{N}$, u_n is an H^2 -solution to (1.1)–(1.3) where the right-hand side of (1.1) is f_n , and if

$$u_n \xrightarrow[n \rightarrow \infty]{C([0, T]; L^2(\Omega))} u \quad \text{and} \quad f_n \xrightarrow[n \rightarrow \infty]{L^1((0, T); L^2(\Omega))} f, \quad (2.4)$$

for any $T > 0$. Sometimes, we shall write (u, f) , (u, U) or (u, U, f) to designate a solution with the obvious meanings.

We recall that under Assumptions 2.1, if $f \in L^1_{\text{loc}}([0, \infty); L^2(\Omega))$, then for any $u_0 \in L^2(\Omega)$, there exists a unique weak solution to (1.1)–(1.3) ([9, Theorem 2.8]).

Theorem 2.4 (Infinite time extinction property). *Let Assumption 2.1 be fulfilled, let $u_0 \in L^2(\Omega)$, let $f \in L^1((0, \infty); L^2(\Omega))$ and let u be the unique weak solution to (1.1)–(1.3). Then,*

$$\lim_{t \nearrow \infty} \|u(t)\|_{L^2(\Omega)} = 0.$$

Proposition 2.5 (Infinite time extinction property). *Let Assumption 2.1 be fulfilled with $m = 1$, let $f \in L^1((0, \infty); L^2(\Omega))$, let $u_0 \in L^2(\Omega)$ and let u be the unique weak solution to (1.1)–(1.3). If $f = 0$ almost everywhere on (T_0, ∞) , for some $T_0 \geq 0$, then*

$$\|u(t)\|_{L^2(\Omega)} \leq \|u(T_0)\|_{L^2(\Omega)} e^{-\operatorname{Re}(ae^{i\theta})(t-T_0)}.$$

for any $t \geq T_0$.

In order to have finite time extinction of the solutions, we make the following assumptions.

Assumption 2.6. Let Assumption 2.1 be fulfilled with $m < 1$, let $f \in L^1_{\text{loc}}([0, \infty); L^2(\Omega))$, let $u_0 \in L^2(\Omega)$ and let u be the unique weak solution u to (1.1)–(1.3). We assume that there exists a finite time $T_0 \geq 0$ such that

$$\text{for almost every } (t, x) \in (T_0, \infty) \times \Omega, f(t, x) = 0. \quad (2.5)$$

If $m = 0$, then we may make a weaker hypothesis. Instead of (2.5), we may assume that

$$f \in L^\infty((T_0, \infty); L^\infty(\Omega)) \quad \text{and} \quad \|f\|_{L^\infty((T_0, \infty); L^\infty(\Omega))} < \operatorname{Re}(ae^{i\theta}). \quad (2.6)$$

Finally, we set $\delta = \frac{(N+2)-m(N-2)}{N(1-m)+4} \in (\frac{1}{2}, 1)$ and $\lambda = 2(1-\delta) = \frac{4(1-m)}{N(1-m)+4}$.

Theorem 2.7 (Finite time extinction property). *Let Assumption 2.6 be fulfilled.*

1. For any $t \geq T_0$,

$$\|u(t)\|_{L^2(\Omega)} \leq \left(\|u(T_0)\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} - \lambda M C_{\text{GN}}^{-\frac{4}{N(1-m)+4}} (t-T_0) \right)_+^{\frac{N(1-m)+4}{4(1-m)}}, \quad (2.7)$$

where C_{GN} is given by (3.18) below and

$$M = \min \{ \cos \theta, \operatorname{Re}(ae^{i\theta}) - \|f\|_{L^\infty((T_0, \infty); L^\infty(\Omega))} \}. \quad (2.8)$$

In particular,

$$\forall t \geq T_\star, \|u(t)\|_{L^2(\Omega)} = 0, \quad (2.9)$$

where

$$T_\star \leq \frac{C_{\text{GN}}^{\frac{4}{N(1-m)+4}}}{\lambda M} \|u(T_0)\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} + T_0. \quad (2.10)$$

2. There exists $\varepsilon_\star = \varepsilon_\star(m, N)$ satisfying the following property. If

$$\begin{cases} \|u_0\|_{L^2(\Omega)}^{2(1-\delta)} \leq \varepsilon_\star T_0, \\ \|f(t)\|_{L^2(\Omega)}^2 \leq \varepsilon_\star (T_0 - t)^{\frac{2\delta-1}{1-\delta}}, \end{cases} \quad (2.11)$$

for almost every $t > 0$, then (2.9) holds true with $T_\star = T_0$.

Remark 2.8. Here are some comments about Theorem 2.7.

1. If f satisfies (2.5), then $\|f\|_{L^\infty((T_0, \infty); L^\infty(\Omega))} = 0$ and (2.8) reads as

$$M = \min \{ \cos \theta, \operatorname{Re}(ae^{i\theta}) \}.$$

2. We have that $\frac{2\delta-1}{1-\delta} = \frac{N(1-m)+4m}{2(1-m)}$.

3. PROOF OF THE FINITE TIME EXTINCTION PROPERTY

If u is an H^2 -strong solution, then the map $t \mapsto \|u(t)\|_{L^2(\Omega)}^2$ belongs to $W_{\text{loc}}^{1, \infty}([0, \infty); \mathbb{R})$ and

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u(t)\|_{L^2(\Omega)}^2 + \cos \theta \|\nabla u(t)\|_{L^2(\Omega)}^2 + \operatorname{Re}(ae^{i\theta}) \|u(t)\|_{L^{m+1}(\Omega)}^{m+1} + \operatorname{Re}(be^{i\theta}) \|u(t)\|_{L^{p+1}(\Omega)}^{p+1} \\ & + \operatorname{Re}(\gamma e^{i\theta}) \|u(t)\|_{L^2(\Omega)}^2 = \operatorname{Re} \left(e^{i\theta} \int_{\Omega} f(t, x) \overline{u(t, x)} \, dx \right), \end{aligned} \quad (3.1)$$

for almost every $t > 0$ ([9, Theorem 2.14]). The proof of Property 1 of Theorem 2.7 (as well as Property 2) relies on the estimate of the time derivative of the mass (3.1) to arrive at the estimate

$$y'(t) + \nu y(t)^\delta \leq 0, \quad (3.2)$$

where $y(\cdot) = \|u(\cdot)\|_{L^2(\Omega)}^2$, for some $\nu > 0$ and $\delta \in (0, 1)$. But (3.1) does not hold for the weak solutions, as well as (3.2). As a consequence, we first prove Property 1 for the strong solutions and then proceed by density. The passage to the limit is possible with the help of the continuous dependence (see (3.20) below) and the weak solutions are approached by strong solutions with the help of Lemma 3.1 below. This proves the extinction of the solution in finite time. But the proof of Property 2 of Theorem 2.7, which permits us to choose at which time the solution vanishes, is more delicate. To this end, we use again the estimate of the time derivative of the mass (3.1) and the assumption

$$\|f(t)\|_{L^2(\Omega)}^2 \leq \varepsilon_\star (T_0 - t)_+^{\frac{2\delta-1}{1-\delta}}, \quad (3.3)$$

for almost every $t > 0$. We then obtain (3.2), and we then apply [5, Lemma 5.2] to obtain the extinction of the solution at time T_0 . Again, we have to consider strong solutions. But the key assumption (3.3) cannot be obtained for a smooth sequence $(f_n)_{n \in \mathbb{N}}$ that approaches the external source f . Rather, we first prove a more general result (Lemma 3.2 below) than [5, Lemma 5.2], which permits us to prove Property 2 by density.

Lemma 3.1. *Let I be an interval (not necessarily open) with $-\infty \leq \inf I < \sup I \leq \infty$, let $1 \leq p < \infty$, let X be a Banach space and let $f \in L^p_{\text{loc}}(I; X)$. Then there exist $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}(I; X)$ and $g \in L^p_{\text{loc}}(I; \mathbb{R})$ such that,*

$$f_n \xrightarrow[n \rightarrow \infty]{L^p_{\text{loc}}(I; X)} f, \quad (3.4)$$

$$\text{for a.e. } t \in I, f_n(t) \xrightarrow[n \rightarrow \infty]{X} f(t), \quad (3.5)$$

$$\text{for a.e. } t \in I \text{ and any } n \in \mathbb{N}, \|f_n(t)\|_X \leq g(t), \quad (3.6)$$

$$\text{for any } n \in \mathbb{N}, \text{supp } f_n \subset \text{supp } f + \bar{B}\left(0, \frac{1}{n}\right). \quad (3.7)$$

If, in addition, $f \in L^p(I; X)$, then

$$f_n \xrightarrow[n \rightarrow \infty]{L^p(I; X)} f \text{ and } g \in L^p(I; \mathbb{R}). \quad (3.8)$$

Finally, if for some $q \in [1, \infty]$ and a Banach space Y , $f \in L^q(I; Y)$, then $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}(I; Y)$ and

$$\|f_n\|_{L^q(I; Y)} \leq \|f\|_{L^q(I; Y)}, \quad (3.9)$$

for any $n \in \mathbb{N}$.

Proof. Let $f \in L^p_{\text{loc}}(I; X)$. Let for each $n \in \mathbb{N}$, $I_n = (\inf I + \frac{1}{n}, \sup I - \frac{1}{n})$. Finally, let $(\rho_n)_{n \in \mathbb{N}}$ be a sequence of mollifiers. Let us denote by \tilde{f} the extension of f by 0 outside I . Let $n, j \in \mathbb{N}$. We define $g_{n,j} = \rho_j \star (\tilde{f} \mathbb{1}_{I_n})|_I$. It is well-known that for any $n \in \mathbb{N}$, $(g_{n,j})_{j > n} \subset \mathcal{D}(I; X)$ and $g_{n,j} \xrightarrow[j \rightarrow \infty]{L^p(I; X)} f \mathbb{1}_{I_n}$, as $j \rightarrow \infty$. See, for instance, Droniou [16, Théorème 1.7.1, p.27]. See also Brezis [10] (Proposition 4.18, p.106 and Theorem 4.22, p.109). It follows that there exists an increasing function $\psi : \mathbb{N} \rightarrow \mathbb{N}$ such that for any $n \in \mathbb{N}$, setting $f_n = g_{n, \psi(n)}$, we have $f_n - f \mathbb{1}_{I_n} \xrightarrow[n \rightarrow \infty]{L^p(I; X)} 0$, as $n \rightarrow \infty$. By the partial converse of the dominated convergence theorem for vector-valued functions (Droniou [16, Théorème 1.3.4, p.16]), we may assume, by renumbering the sequence if necessary, that for a.e. $t \in I$, $f_n(t) - f \mathbb{1}_{I_n}(t) \xrightarrow[n \rightarrow \infty]{X} 0$, as $n \rightarrow \infty$, and $\|f_n - f \mathbb{1}_{I_n}\|_X \leq g \in L^p(I; \mathbb{R})$, a.e. in I and for any $n \in \mathbb{N}$. Since for any compact interval $J \subset I$ and $t \in \overset{\circ}{I}$ (the interior of I), there exists $n_0 \in \mathbb{N}$ such that for any $n > n_0$, $J \subset I_n$ and $t \in I_n$, we easily conclude that $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}(I; X)$ satisfies (3.4) and (3.5) (and also (3.8) if $f \in L^p(I; X)$). Since for any $n \in \mathbb{N}$, $\text{supp } \rho_n = \bar{B}(0, \frac{1}{n})$, (3.7) comes from a classical result of the convolution of two functions (Brezis [10, Proposition 4.18]). Finally, if $f \in L^q(I; Y)$ for some $q \in [1, \infty]$ and a Banach space Y , then we have that $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}(I; Y)$ and (3.9) comes from Young's inequality for vector-valued functions (Droniou [16, Proposition 1.7.1, p.25]). \square

Lemma 3.2. *Let $\alpha, \delta > 0$. Let $g \in L^1_{\text{loc}}([0, \infty); \mathbb{R})$ be a nonnegative function and let $t_0 \geq 0$.*

1. *For any $z_0 \geq 0$, there exists a unique solution $z \in W^{1,1}_{\text{loc}}([t_0, \infty); \mathbb{R})$ to*

$$\begin{cases} \forall t \geq t_0, z(t) \geq 0, \\ \text{for a.e. } t > t_0, z'(t) + \alpha z(t)^\delta = g(t), \end{cases} \quad (3.10)$$

such that

$$z(t_0) = z_0. \quad (3.11)$$

Let $g_1, g_2 \in L^1_{\text{loc}}([0, \infty); \mathbb{R})$ be nonnegative functions and let $z_1, z_2 \in W^{1,1}_{\text{loc}}([t_0, \infty); \mathbb{R})$ be solutions to

$$\begin{cases} \forall t \geq t_0, z_j(t) \geq 0, \\ \text{for a.e. } t > 0, z'_j(t) + \alpha z_j(t)^\delta = g_j(t), \end{cases} \quad (3.12)$$

for $j \in \{1, 2\}$. Then,

$$|z_1(t) - z_2(t)| \leq |z_1(s) - z_2(s)| + \int_s^t |g_1(\sigma) - g_2(\sigma)| d\sigma, \quad (3.13)$$

for any $t \geq s \geq t_0$.

2. *Let $z \in W^{1,1}_{\text{loc}}([t_0, \infty); \mathbb{R})$ be a solution to (3.10) and let $y \in W^{1,1}_{\text{loc}}([t_0, \infty); \mathbb{R})$ be a nonnegative solution to*

$$\text{for a.e. } t > t_0, y'(t) + \alpha y(t)^\delta \leq g(t), \quad (3.14)$$

If for some $t_\star \in [t_0, T)$, $y(t_\star) \leq z(t_\star)$, then

$$\forall t \geq t_\star, y(t) \leq z(t). \quad (3.15)$$

Proof. Let $\alpha, \delta > 0$. Let $g \in L^1_{\text{loc}}([0, \infty); \mathbb{R})$ with $g \geq 0$, a.e. in $(0, \infty)$, and let $t_0 \geq 0$.

Proof of Property 1. Let $([t_0, T_{\text{max}}), z)$ be any maximal solution to (3.10) with $T_{\text{max}} < \infty$. Then for any $t \in [t_0, T_{\text{max}})$,

$$\forall t \in [t_0, T_{\text{max}}), 0 \leq z(t) \leq z(t_0) + \int_0^{T_{\text{max}}} g(s) ds < \infty. \quad (3.16)$$

Now, let $z_0 \geq 0$ and set for a.e. $t > 0$ and any $x \in \mathbb{R}$, $f(t, x) = g(t) - \alpha (x \mathbb{1}_{[0, \infty)}(x))^\delta$. Then by Carathéodory's Theorem and Zorn's Lemma, there exist $t_0 < T_{\text{max}} \leq \infty$ and a maximal solution $z \in W^{1,1}_{\text{loc}}([t_0, T_{\text{max}}); \mathbb{R})$ to $z' = f(\cdot, z)$, a.e. on (t_0, T_{max}) , that is

$$\text{for a.e. } t \in (t_0, T_{\text{max}}), z'(t) + \alpha (z(t) \mathbb{1}_{\{z(t) \geq 0\}}(t))^\delta = g(t), \quad (3.17)$$

such that $z(t_0) = z_0$. In addition, the following blow-up alternative holds true: if $T_{\max} < \infty$, then $\lim_{t \nearrow T_{\max}} |z(t)| = \infty$. Now, assume by contradiction that for some $t_1 \in (t_0, T_{\max})$, $z(t_1) < 0$. Then, since $z(t_0) \geq 0$, we obtain by continuity the existence of a $T \in [t_0, T_{\max})$ and of a $\delta \in (0, T_{\max} - T)$ such that $z(T) = 0$ and for any $t \in (T, T + \delta]$, $z(t) < 0$. It then follows from (3.17) that $z' \geq 0$, a.e. on $(T, T + \delta)$, so that $0 = z(T) \leq z(T + \delta) < 0$, a contradiction. It follows that,

$$\forall t \in [t_0, T_{\max}), z(t) \geq 0,$$

and by (3.16) and the blow-up alternate, we obtain $T_{\max} = \infty$. As a consequence, any maximal solution to (3.10) is global. Now, let g_1, g_2 and z_1, z_2 be as in the statement of the lemma. Let $z = z_1 - z_2$ and $g = g_1 - g_2$. It follows that

$$\text{for a.e. } t > t_0, z'(t) + \alpha(z_1(t)^\delta - z_2(t)^\delta) = g(t).$$

Multiplying by z , using that $s \mapsto \alpha s^\delta$ is increasing over $[0, \infty)$ and integrating, we get that z satisfies,

$$\forall t \geq t_0 |z_1(t) - z_2(t)| \leq |z_1(s) - z_2(s)| + \int_s^t |g_1(\sigma) - g_2(\sigma)| d\sigma.$$

In particular, this implies uniqueness of the solution, and Property 1 is proved.

Proof of Property 2. Let the assumptions be fulfilled. If (3.15) does not hold, then since $y(t_*) \leq z(t_*)$, we have by continuity that there exist $t_* \leq T_* < \infty$ and $\varepsilon > 0$ such that $y(T_*) = z(T_*)$ and $y(t) > z(t)$, for any $t \in (T_*, T_* + \varepsilon)$. This leads with (3.10) and (3.14) to $y' \leq z'$, almost everywhere on $(T_*, T_* + \varepsilon)$. Integrating over (T_*, t) for $t \in (T_*, T_* + \varepsilon)$, we obtain that $y(t) \leq z(t)$, for any $t \in [T_*, T_* + \varepsilon]$, a contradiction. Hence, the result. \square

Let us recall the following Gagliardo–Nirenberg inequality (Gagliardo [18], Nirenberg [19]). Let Ω be an open subset of \mathbb{R}^N and let $0 \leq m \leq 1$. Then there exists $C_{\text{GN}} = C_{\text{GN}}(m, N)$ such that for any $u \in H_0^1(\Omega) \cap L^{m+1}(\Omega)$,

$$\|u\|_{L^2(\Omega)}^{\frac{(N+2)-m(N-2)}{2}} \leq C_{\text{GN}} \|u\|_{L^{m+1}(\Omega)}^{m+1} \|\nabla u\|_{L^2(\Omega)}^{\frac{N(1-m)}{2}}. \quad (3.18)$$

It follows that

$$\begin{aligned} & \|u\|_{L^2(\Omega)}^{\frac{(N+2)-m(N-2)}{2}} \\ & \leq C_{\text{GN}} \left(\|\nabla u\|_{L^2(\Omega)}^2 + \|u\|_{L^{m+1}(\Omega)}^{m+1} \right) \left(\|\nabla u\|_{L^2(\Omega)}^2 + \|u\|_{L^{m+1}(\Omega)}^{m+1} \right)^{\frac{N(1-m)}{4}} \\ & = C_{\text{GN}} \left(\|\nabla u\|_{L^2(\Omega)}^2 + \|u\|_{L^{m+1}(\Omega)}^{m+1} \right)^{\frac{N(1-m)+4}{4}}, \end{aligned}$$

and then

$$\|u\|_{L^2(\Omega)}^{2\delta} \leq C_{\text{GN}}^{\frac{4}{N(1-m)+4}} \left(\|\nabla u\|_{L^2(\Omega)}^2 + \|u\|_{L^{m+1}(\Omega)}^{m+1} \right), \quad (3.19)$$

for any $u \in H_0^1(\Omega) \cap L^{m+1}(\Omega)$, where δ is defined in Assumption 2.6. Finally, let us recall that if (u, f) and (\tilde{u}, \tilde{f}) are (strong or weak) solutions to (1.1)–(1.2), then

$$\|u(t) - \tilde{u}(t)\|_{L^2(\Omega)} \leq \|u(s) - \tilde{u}(s)\|_{L^2(\Omega)} + \int_s^t \|f(\sigma) - \tilde{f}(\sigma)\|_{L^2(\Omega)} d\sigma, \quad (3.20)$$

for any $t \geq s \geq 0$ ([9, Proposition 2.6]). Now, we are able to prove Theorem 2.7.

Proof of Theorem 2.7. Let f, u_0, u and M be as in the statement of the theorem and set for any $t \geq 0, y(t) = \|u(t)\|_{L^2(\Omega)}^2$.

Proof of Property 1. We only show that u satisfies (2.7), from which (2.9) and (2.10) will follow. We first assume that $f \in \mathcal{D}((0, \infty); L^2(\Omega))$ and $u_0 \in \mathcal{D}(\Omega)$, so that u is an H^2 -solution ([9, Theorem 2.14]). We have by (3.1), (3.19), (2.5) and (2.6) that for a.e. $t > T_0$,

$$y'(t) + 2\alpha y(t)^\delta \leq 0, \quad (3.21)$$

where $\alpha = MC_{\text{GN}}^{-\frac{4}{N(1-m)+4}}$. After integration, we obtain that for any $t \geq T_0$,

$$\|u(t)\|_{L^2(\Omega)}^2 \leq \left(\|u(T_0)\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} - \lambda MC_{\text{GN}}^{-\frac{4}{N(1-m)+4}} (t - T_0) \right)^{\frac{N(1-m)+4}{2(1-m)}}, \quad (3.22)$$

which is (2.7).

End of the proof when $m > 0$. Now, we consider the general case: $u_0 \in L^2(\Omega)$ and $f \in L^1((0, \infty); L^2(\Omega))$ which satisfies (2.5). We apply Lemma 3.1: let $(\varphi_n)_{n \in \mathbb{N}} \subset \mathcal{D}(\Omega)$ and $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}((0, \infty); L^2(\Omega))$ be such that,

$$\varphi_n \xrightarrow[n \rightarrow \infty]{L^2(\Omega)} u_0 \quad \text{and} \quad f_n \xrightarrow[n \rightarrow \infty]{L^1((0, \infty); L^2(\Omega))} f. \quad (3.23)$$

Finally, for each $n \in \mathbb{N}$, let (u_n, f_n) be the H^2 -solution to (1.1)–(1.2) such that $u_n(0) = \varphi_n$. By (2.5), (3.7) and (3.22), we get that for any $n \in \mathbb{N}$ and $t \geq T_0 + \frac{1}{n}$,

$$\|u_n(t)\|_{L^2(\Omega)} \leq \left(\left\| u_n \left(T_0 + \frac{1}{n} \right) \right\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} - \lambda MC_{\text{GN}}^{-\frac{4}{N(1-m)+4}} \left(t - T_0 - \frac{1}{n} \right) \right)^{\frac{N(1-m)+4}{4(1-m)}}. \quad (3.24)$$

By (3.23) and (3.20), we may pass to the limit in (3.24), so that u satisfies (2.7).

End of the proof when $m = 0$. Assume that $u_0 \in L^2(\Omega)$ and $f \in L^1_{\text{loc}}([0, \infty); L^2(\Omega))$ which satisfies (2.6). By Lemma 3.1, there exist $(\varphi_n)_{n \in \mathbb{N}} \subset \mathcal{D}(\Omega)$ and $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}((0, \infty); L^2(\Omega))$ such that,

$$\varphi_n \xrightarrow[n \rightarrow \infty]{L^2(\Omega)} u(T_0) \quad \text{and for any } T > T_0, \quad f_n \xrightarrow[n \rightarrow \infty]{L^1((T_0, T); L^2(\Omega))} f, \quad (3.25)$$

$$\text{for any } n \in \mathbb{N}, \|f_n\|_{L^\infty((T_0, \infty); L^\infty(\Omega))} \leq \|f\|_{L^\infty((T_0, \infty); L^\infty(\Omega))}. \quad (3.26)$$

Let $n \in \mathbb{N}$. Let (u_n, f_n) be the H^2 -solution to (1.1)–(1.2) such that $u_n(T_0) = \varphi_n$ (which is possible by uniqueness of the solution and the invariance of (1.1) by time translation). By (3.26), each f_n satisfies (2.6). Then by (3.22), we have for any $n \in \mathbb{N}$ and $t \geq T_0$,

$$\|u_n(t)\|_{L^2(\Omega)}^2 \leq \left(\|u_n(T_0)\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} - \lambda M_n C_{\text{GN}}^{-\frac{4}{N(1-m)+4}} (t - T_0) \right)_+^{\frac{N(1-m)+4}{2(1-m)}},$$

where

$$M_n = \min \{ \cos \theta, \operatorname{Re}(ae^{i\theta}) - \|f_n\|_{L^\infty((T_0, \infty); L^\infty(\Omega))} \}.$$

By (3.26), $M \leq M_n$, so that

$$\|u_n(t)\|_{L^2(\Omega)}^2 \leq \left(\|u_n(T_0)\|_{L^2(\Omega)}^{\frac{4(1-m)}{N(1-m)+4}} - \lambda M C_{\text{GN}}^{-\frac{4}{N(1-m)+4}} (t - T_0) \right)_+^{\frac{N(1-m)+4}{2(1-m)}}, \quad (3.27)$$

for any $n \in \mathbb{N}$ and $t \geq T_0$. By (3.25) and (3.20), we may pass to the limit in (3.27) and then u satisfies (2.7).

Proof of Property 2. We first note by (2.11) that $f \in L^p((0, \infty); L^2(\Omega))$, where $p = \frac{2\delta}{2\delta-1} > 1$. By Lemma 3.1, there exist $h \in L^p((0, \infty); \mathbb{R})$, $(\varphi_n)_{n \in \mathbb{N}} \subset \mathcal{D}(\Omega)$ and $(f_n)_{n \in \mathbb{N}} \subset \mathcal{D}((0, \infty); L^2(\Omega))$ such that,

$$\varphi_n \xrightarrow[n \rightarrow \infty]{L^2(\Omega)} u_0 \quad \text{and} \quad f_n \xrightarrow[n \rightarrow \infty]{L^p((0, \infty); L^2(\Omega))} f, \quad (3.28)$$

$$\text{for a.e. } t > 0 \quad f_n(t) \xrightarrow[n \rightarrow \infty]{L^2(\Omega)} f(t), \quad (3.29)$$

$$\text{for a.e. } t > 0 \text{ and any } n \in \mathbb{N}, \quad \|f_n(t)\|_{L^2(\Omega)} \leq h(t). \quad (3.30)$$

For each $n \in \mathbb{N}$, let (u_n, f_n) be the H^2 -solution to (1.1)–(1.2) such that $u_n(0) = \varphi_n$, and set for any $t \geq 0$, $y_n(t) = \|u_n(t)\|_{L^2(\Omega)}^2$. Let $n \in \mathbb{N}$. We have by (3.1), (3.19) and Cauchy–Schwarz' inequality that for a.e. $t > 0$,

$$y_n'(t) + 2\alpha y_n(t)^\delta \leq 2 \|f_n(t)\|_{L^2(\Omega)} y_n(t)^{\frac{1}{2}}, \quad (3.31)$$

where $\alpha = \min \{ \cos \theta, \operatorname{Re}(ae^{i\theta}) \} C_{\text{GN}}^{-\frac{4}{N(1-m)+4}}$. Now, we set

$$\varepsilon_* = \min \left\{ (2\delta - 1)^{-\frac{2\delta-1}{\delta}} (\alpha\delta)^{\frac{1}{1-\delta}} (1 - \delta)^{\frac{2\delta-1}{\delta(1-\delta)}}, \alpha\delta(1 - \delta) \right\}.$$

Applying Young's inequality to (3.31) we arrive at,

$$y_n'(t) + 2\alpha y_n(t)^\delta \leq \frac{2\delta - 1}{\delta} (\alpha\delta)^{-\frac{1}{2\delta-1}} \|f_n(t)\|_{L^2(\Omega)}^{\frac{2\delta}{2\delta-1}} + \alpha y_n(t)^\delta,$$

for a.e. $t > 0$, and then

$$\text{for a.e. } t > 0, \quad y_n'(t) + \alpha y_n(t)^\delta \leq g_n(t),$$

where $g_n(t) = \frac{2\delta-1}{\delta}(\alpha\delta)^{-\frac{1}{2\delta-1}}\|f_n(t)\|_{L^2(\Omega)}^{\frac{2\delta}{2\delta-1}}$. Let $z_n \in W_{\text{loc}}^{1,1}([0, \infty); \mathbb{R})$, with $z_n \geq 0$ everywhere in $[0, \infty)$, be the unique solution to

$$\text{for a.e. } t > 0, \quad z'_n(t) + \alpha z_n(t)^\delta = g_n(t),$$

such that $z_n(0) = y_n(0)$. By Lemma 3.2, we have for any $t \geq 0$, $y_n(t) \leq z_n(t)$. By (3.29), (3.30) and the dominated convergence theorem, we have that $g_n \xrightarrow[n \rightarrow \infty]{L^1((0, \infty); \mathbb{R})} g$, where for a.e. $t > 0$,

$$g(t) = \frac{2\delta-1}{\delta}(\alpha\delta)^{-\frac{1}{2\delta-1}}\|f(t)\|_{L^2(\Omega)}^{\frac{2\delta}{2\delta-1}}.$$

We then infer with the help of (3.28), (3.20) and Lemma 3.2 that

$$\forall t \geq 0, \quad y(t) \leq z(t), \tag{3.32}$$

where $z \in W_{\text{loc}}^{1,1}([0, \infty); \mathbb{R})$ is the unique nonnegative solution to

$$\text{for a.e. } t > 0, \quad z'(t) + \alpha z(t)^\delta = g(t),$$

such that $z(0) = y(0)$. By (2.11), it follows that

$$\text{for a.e. } t > 0, \quad z'(t) + \alpha z(t)^\delta \leq z_*(T_0 - t)_+^{\frac{\delta}{1-\delta}},$$

where $z_* = (\alpha\delta^\delta(1-\delta))^{\frac{1}{1-\delta}}$. Finally, let $\zeta_* = (\alpha\delta(1-\delta)T_0)^{\frac{1}{1-\delta}}$, and let $\zeta \in W_{\text{loc}}^{1,1}([0, \infty); \mathbb{R})$, with $\zeta \geq 0$ everywhere in $[0, \infty)$, be the unique solution to

$$\text{for a.e. } t > 0, \quad \zeta'(t) + \alpha\zeta(t)^\delta = z_*(T_0 - t)_+^{\frac{\delta}{1-\delta}},$$

such that $\zeta(0) = \zeta_*$. By (2.11), we have $z(0) \leq \zeta_*$ and then Lemma 3.2 implies that

$$\forall t \geq 0, \quad z(t) \leq \zeta(t), \tag{3.33}$$

Finally, by the uniqueness of the solution, we obtain

$$\forall t \geq 0, \quad \zeta(t) = \zeta_* T_0^{-\frac{1}{1-\delta}} (T_0 - t)_+^{\frac{1}{1-\delta}}. \tag{3.34}$$

Putting together (3.32)–(3.34), we get that

$$\forall t \geq 0, \quad y(t) \leq \zeta_* T_0^{-\frac{1}{1-\delta}} (T_0 - t)_+^{\frac{1}{1-\delta}},$$

from which the result follows. \square

Proof of Proposition 2.5. By density (in particular (3.7)) and continuous dependence (3.20), we may assume $f \in \mathcal{D}((0, \infty); L^2(\Omega))$, $u_0 \in \mathcal{D}(\Omega)$ so that u is an H^2 -solution ([9, Theorem 2.14]). By (3.1), we have

$$\forall t \geq T_0, \quad \frac{1}{2} \frac{d}{dt} \|u(t)\|_{L^2(\Omega)}^2 + \text{Re}(ae^{i\theta}) \|u(t)\|_{L^2(\Omega)}^2 \leq 0,$$

from which the result follows. \square

Proof of Theorem 2.4. By (3.20), density and Proposition 2.5, we may assume that $u_0 \in \mathcal{D}(\Omega)$, $f \in \mathcal{D}((0, \infty); L^2(\Omega))$ and $m < 1$. The result then comes from Theorem 2.7. \square

Acknowledgements

Pascal Bégout acknowledges funding from ANR under grant ANR-17-EUR-0010 (Investissements d’Avenir program). The research of J. I. Díaz was partially supported by the project PID-2020-112517GBI00 of the AEI and MCIU/AEI/10.13039/-501100011033/FEDER, EU.


REFERENCES

- [1] S. Antontsev, J.-P. Dias, M. Figueira, *Complex Ginzburg–Landau equation with absorption: existence, uniqueness and localization properties*, J. Math. Fluid Mech. **16** (2014), no. 2, 211–223.
- [2] S.N. Antontsev, J.I. Díaz, S. Shmarev, *Energy Methods for Free Boundary Problems*, Progress in Nonlinear Differential Equations and Their Applications, 48, Birkhäuser, Boston, MA, 2002.
- [3] P. Bégout, *Finite time extinction for a damped nonlinear Schrödinger equation in the whole space*, Electron. J. Differential Equations (2020), no. 39, 1–18.
- [4] P. Bégout, *The dual space of a complex Banach space restricted to the field of real numbers*, Adv. Math. Sci. Appl. **31** (2022), no. 2, 241–252.
- [5] P. Bégout, J.I. Díaz, *Finite time extinction for the strongly damped nonlinear Schrödinger equation in bounded domains*, J. Differential Equations **268** (2020), no. 7, 4029–4058.
- [6] P. Bégout, J.I. Díaz, *Finite time extinction for a class of damped Schrödinger equations with a singular saturated nonlinearity*, J. Differential Equations **308** (2022), 252–285.
- [7] P. Bégout, J.I. Díaz, *Finite time extinction for a critically damped Schrödinger equation with a sublinear nonlinearity*, Adv. Differential Equations **28** (2023), no. 3–4, 311–340.
- [8] P. Bégout, J.I. Díaz, *Strong stabilization of damped nonlinear Schrödinger equation with saturation on unbounded domains*, J. Math. Anal. Appl. **538** (2024), 128329.
- [9] P. Bégout, J.I. Díaz, *Damped nonlinear Ginzburg–Landau equation with saturation. Part I. Existence of solutions on general domains*, Opuscula Math. **46** (2026), no. 2, 153–183.
- [10] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, New York, 2011.
- [11] A.C. Casal, J.I. Díaz, *On the principle of pseudo-linearized stability: applications to some delayed nonlinear parabolic equations*, Nonlinear Anal. **63** (2005), e997–e1007.
- [12] A.C. Casal, J.I. Díaz, *On the complex Ginzburg–Landau equation with a delayed feedback*, Math. Models Methods Appl. Sci. **16** (2006), no. 1, 1–17.
- [13] A.C. Casal, J.I. Díaz, M. Stich, *On some delayed nonlinear parabolic equations modeling CO oxidation*, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. **13B** (2006), 413–426.

- [14] A.C. Casal, J.I. Díaz, M. Stich, *Control of turbulence in oscillatory reaction-diffusion systems through a combination of global and local feedback*, Phys. Rev. E **76** (2007), 036209.
- [15] J.I. Díaz, J.F. Padiál, J.I. Tello, L. Tello, *Complex Ginzburg–Landau equations with a delayed nonlocal perturbation*, Electron. J. Differential Equations (2020), Paper no. 40.
- [16] J. Droniou, *Intégration et Espaces de Sobolev à Valeurs Vectorielles*, HAL preprint hal-01382368, 2001.
- [17] K. Fenza, M. Labbadi, M. Ouzahra, *Finite-time stabilization of a class of nonlinear systems in Hilbert space*, [in:] Proc. 2025 IEEE 64th Conference on Decision and Control (CDC), IEEE, 2025, 3003–3008.
- [18] E. Gagliardo, *Ulteriori proprietà di alcune classi di funzioni in più variabili*, Ricerche Mat. **8** (1959), 24–51.
- [19] L. Nirenberg, *On elliptic partial differential equations*, Ann. Scuola Norm. Sup. Pisa (3) **13** (1959), 115–162.
- [20] L. Rosier, B.-Y. Zhang, *Null controllability of the complex Ginzburg–Landau equation*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **26** (2009), no. 2, 649–673.

Pascal Bégout (corresponding author)


Pascal.Begout@math.cnrs.fr

 <https://orcid.org/0000-0002-9172-3057>

Toulouse School of Economics
Université Toulouse Capitole
Institut de Mathématiques de Toulouse
1, Esplanade de l'Université
31080 Toulouse Cedex 6, France

Jesús Ildefonso Díaz

jidiaz@ucm.es

 <https://orcid.org/0000-0003-1730-9509>

Universidad Complutense de Madrid
Instituto de Matemática Interdisciplinar
Plaza de las Ciencias, 3
28040 Madrid, Spain

Received: February 4, 2026.

Revised: March 10, 2026.

Accepted: March 11, 2026.

Published online: April 10, 2026.